

- Holt, H.E., Francis, I.L., Field, J., Herderich, M.J. & Iland, P.G., 2008. Relationships between berry size, berry phenolic composition and wine quality scores for Cabernet Sauvignon (*Vitis vinifera* L.) from different pruning treatments and different vintages. *Aust. J. Grape Wine Res.* 14(3), 191-202.
- Houel, C., Martin-Magniette, M.L., Nicolas, S.D., Lacombe, T., Le-Cunff, L., Franck, D., Torregrosa, L., Conéjéro, G., Lalet, S., This, P. & Adam-Blondon, A.F., 2013. Genetic variability of berry size in the grapevine (*Vitis vinifera* L.). *Aust. J. Grape Wine Res.* 19(2), 208-220.
- Kalua, C.M. & Boss, P.K., 2009. Evolution of volatile compounds during the development of Cabernet Sauvignon grapes (*Vitis vinifera* L.). *J. Agric. Food Chem.* 57(9), 3818-3830.
- Keller, M., 2010 (1st ed.). *The science of grapevines: Anatomy and physiology*. Elsevier Academic Press, Burlington, MA.
- Keller, M., Mills, L.J. & Harbertson, J.F., 2011. Rootstock effects on deficit-irrigated winegrapes in a dry climate: Vigor, yield formation, and fruit ripening. *Am. J. Enol. Vitic.* 63(1), 29-39.
- Kennedy, J.A., Matthews, M.A. & Waterhouse, A.L., 2002. Effect of maturity and vine water status on grape skin and wine flavonoids. *Am. J. Enol. Vitic.* 53(4), 268-274.
- Kontoudakis, N., Esteruelas, M., Fort, F., Canals, J.M., De Freitas, V. & Zamora, F., 2011. Influence of the heterogeneity of grape phenolic maturity on wine composition and quality. *Food Chem.* 124(3), 767-774.
- Matthews, M.A. & Anderson, M.M., 1988. Fruit ripening in *Vitis vinifera* L.: Response to seasonal water deficits. *Am. J. Enol. Vitic.* 39(4), 313-320.
- Melo, M.S., Schultz, H.R., Volschenk, C.G. & Hunter, J.J., 2015. Berry size variation of *Vitis vinifera* L. cv. Syrah: Morphological dimensions, berry composition and wine quality. *S. Afr. J. Enol. Vitic.* 36(1), 1-10.
- Ojeda, H., Andary, C., Kraeva, E., Carbonneau, A. & Deloire, A., 2002. Influence of pre- and postveraison water deficit on synthesis and concentration of skin phenolic compounds during berry growth of *Vitis vinifera* cv. Shiraz. *Am. J. Enol. Vitic.* 53(4), 261-267.
- Peinado, R.A., Moreno, J., Bueno, J.E., Moreno, J.A. & Mauricio, J.C., 2004. Comparative study of aromatic compounds in two young white wines subjected to pre-fermentative cryomaceration. *Food Chem.* 84(4), 585-590.
- Pérez-Magariño, S. & González-Sanjosé, M.L., 2003. Application of absorbance values used in wineries for estimating CIELAB parameters in red wines. *Food Chem.* 81(2), 301-306.
- Petrie, P.R. & Clingeleffer, P.R., 2006. Crop thinning (hand versus mechanical), grape maturity and anthocyanin concentration: Outcomes from irrigated Cabernet Sauvignon (*Vitis vinifera* L.) in a warm climate. *Aust. J. Grape Wine Res.* 12(1), 21-29.
- Pisciotta, A., Di Lorenzo, R., Barbagallo, M.G. & Hunter, J.J., 2013. Berry characterisation of cv. Shiraz according to position on the rachis. *S. Afr. J. Enol. Vitic.* 34(1), 100-107.
- Ribéreau-Gayon, P., Glories, Y., Maujean, A. & Dubourdieu, D., 2006. Phenolic compounds. In *Handbook of enology*, vol. 2. Wiley, Chichester.
- Robinson, J., 2006 (3rd ed.). *The Oxford companion to wine*. Oxford, Oxford University Press.
- Roby, G. & Matthews, M., 2004. Relative proportions of seed, skin and flesh, in ripe berries from Cabernet Sauvignon grapevines grown in a vineyard either well irrigated or under water deficit. *Aust. J. Grape Wine Res.* 10(1), 74-82.
- Roby, G., Harbertson, J.F., Adams, D.A. & Matthews, M.A., 2004. Berry size and vine water deficits as factors in winegrape composition: anthocyanins and tannins. *Aust. J. Grape Wine Res.* 10(2), 100-107.
- Rolle, L., Torchio, F., Giacosa, S. & Río Segade, S., 2015. Berry density and size as factors related to the physicochemical characteristics of Muscat Hamburg table grapes (*Vitis vinifera* L.). *Food Chem.* 173, 105-113.
- Rolle, L., Torchio, F., Giacosa, S., Río Segade, S., Cagnasso, E. & Gerbi, V., 2012. Assessment of physicochemical differences in Nebbiolo grape berries from different production areas and sorted by flotation. *Am. J. Enol. Vitic.* 62(2), 195-204.
- Sabir, A., Kafkas, E. & Tangolar, S., 2010. Distribution of major sugars, acids, and total phenols in juice of five grapevine (*Vitis* spp.) cultivars at different stages of berry development. *Span. J. Agric. Res.* 8(2), 425-433.
- Selli, S., Cabaroglu, T., Canbas, A., Erten, H. & Nurgel, C., 2003. Effect of skin contact on the aroma composition of the musts of *Vitis vinifera* L. cv. Muscat of Bornova and Narince grown in Turkey. *Food Chem.* 81(3), 341-347.
- Shellie, K.C., 2010. Water deficit effect on ratio of seed to berry fresh weight and berry weight uniformity in winegrape cv. Merlot. *Am. J. Enol. Vitic.* 61(3), 414-418.
- Singleton, V.L., 1972. Effects on red wine composition of removing juice before fermentation to simulate variation in berry size. *Am. J. Enol. Vitic.* 23(3), 106-113.
- Singleton, V.L. & Rossi, J.A., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16, 144-158.
- Smart, R.E., Dick, J.K., Gravett, I.M. & Fisher, B.M., 2017. Canopy management to improve grape yield and wine quality-principles and practices. *S. Afr. J. Enol. Vitic.* 11(1), 3-17.
- Šuklje, K., Lisjak, K., Baša Česnik, H., Janež, L., Du Toit, W., Coetzee, Z., Vanzo, A. & Deloire, A., 2012. Classification of grape berries according to diameter and total soluble solids to study the effect of light and temperature on methoxypyrazine, glutathione, and hydroxycinnamate evolution during ripening of Sauvignon Blanc (*Vitis vinifera* L.). *J. Agric. Food Chem.* 60(37), 9454-9461.
- Swiegers, J.H. & Pretorius, I.S., 2005. Yeast modulation of wine flavor. *Adv. Appl. Microbiol.* 57, 131-175.
- Walker, R.R., Blackmore, D.H., Clingeleffer, P.R., Kerridge, G.H., Rühl, E.H. & Nicholas, P.R., 2005. Shiraz berry size in relation to seed number and implications for juice and wine composition. *Aust. J. Grape Wine Res.* 11(1), 2-8.
- Wen, Y.Q., Cui, J., Zhang, Y., Duan, C.Q. & Pan, Q.H., 2014. Comparison of organic acid levels and *L*-IdnDH expression in Chinese-type and European-type grapes. *Euphytica* 196(1), 63-76.
- Wong, D.C.J., Gutierrez, R.L., Dimopoulos, N., Gambetta, G.A. & Castellari, S.D., 2016. Combined physiological, transcriptome, and cis-regulatory element analyses indicate that key aspects of ripening, metabolism, and transcriptional program in grapes (*Vitis vinifera* L.) are differentially modulated accordingly to fruit size. *BMC Genomics* 17(1), 416.
- Zhang, M.X., Pan, Q.H., Yan, G.L. & Duan, C.Q., 2011. Using headspace solid phase micro-extraction for analysis of aromatic compounds during alcoholic fermentation of red wine. *Food Chem.* 125(2), 743-749.