

## Biological survey of the Prince Edward Islands, December 2008

J. Cooper<sup>a,b\*</sup>, M.N. Bester<sup>c</sup>, S.L. Chown<sup>b</sup>,  
R.J.M. Crawford<sup>a,d</sup>, R. Daly<sup>e</sup>, E. Heyns<sup>e</sup>,  
T. Lamont<sup>d</sup>, P.G. Ryan<sup>f</sup> and J.D. Shaw<sup>b</sup>

---

**A biological survey of the Prince Edward Islands took place in December 2008. The survey repeated an earlier survey of the populations of surface-nesting seabirds on both islands and of fur seals (*Arctocephalus* spp.) and alien plants on Prince Edward Island in December 2001. Observations on burrowing seabirds, macro-invertebrates and plant communities on Prince Edward Island and an oceanographic survey of surrounding waters were also included. The survey confirmed many of the observations made on the earlier survey and permitted an assessment of trends in the abundance and distribution of biota since 2001.**

**Key words:** Prince Edward Islands, sub-Antarctic, Southern Ocean, population surveys, alien plants, oceanographic research

---

The Prince Edward Islands (comprising Marion Island and Prince Edward Island) in the southern Indian Ocean are South Africa's only overseas possession and form a Special Nature Reserve with the highest level of national legal protection, equivalent to World Conservation Union (IUCN) Management Category Ia: Strict Nature Reserve (protected area set aside to protect biodiversity and managed mainly for science).<sup>1</sup> In 2007 the islands (including waters to 500 m offshore) were designated a Site of International Importance under the Ramsar Convention on Wetlands ([http://www.ramsar.org/profile/profiles\\_south\\_africa.htm](http://www.ramsar.org/profile/profiles_south_africa.htm)). South Africa also is party to two international instruments that have specific relevance to the Prince Edward Islands: the Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR; [www.ccamlr.org](http://www.ccamlr.org)) and the Agreement on the Conservation of Albatrosses and Petrels (ACAP; [www.acap.aq](http://www.acap.aq)). South Africa has contributed information on penguins to CCAMLR's Ecosystem Monitoring Program since 1994.<sup>2,3</sup> Its ratification of ACAP in November 2003 brought this agreement into force on 1 February 2004. In August 2008, South Africa hosted meetings of ACAP's Working Groups and its Advisory Committee. Nine of the 26 species of albatrosses and petrels listed on ACAP's Annex 1 breed regularly on the Prince Edward Islands.

Three major conservation initiatives for the island group are currently under consideration by the South African authorities: the formal adoption of a new environmental management plan

<sup>a</sup>Animal Demography Unit, Department of Zoology, University of Cape Town, Rondebosch 7701, South Africa.

<sup>b</sup>DST/NRF Centre of Excellence for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa.

<sup>c</sup>Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria 0002, South Africa.

<sup>d</sup>Marine & Coastal Management, Department of Agriculture, Forestry and Fisheries, Private Bag X2, Roggebaai 8012, South Africa.

<sup>e</sup>Southern Ocean Group, Department of Zoology and Entomology, Rhodes University, P.O. Box 94, Grahamstown 6140, South Africa.

<sup>f</sup>Percy FitzPatrick Institute DST/NRF Centre of Excellence, University of Cape Town, Rondebosch 7701, South Africa.

\*Author for correspondence E-mail: john.cooper@uct.ac.za

drafted in 2006 to replace the current plan adopted in 1996; the declaration of a large Marine Protected Area, including all territorial waters out to 12 nautical miles (nm) and extending in several directions to the limit of the islands' 200-nm Exclusive Economic Zone; and the nomination of the islands as a Natural Site to the World Heritage Convention<sup>4-8</sup> (see also <http://whc.unesco.org/en/tentativelists/1923>).

In the main, biological research conducted on the Prince Edward Islands has been restricted to the larger Marion Island, which supports a combined weather and research station, relieved annually.<sup>9</sup> Visits to uninhabited Prince Edward Island occur more rarely and normally only during the annual relief voyages to Marion Island in April/May. Such visits are subject to strict controls on party size, duration and interval.<sup>4,5</sup> In addition, most physical, chemical and biological oceanographic research conducted in the vicinity of the islands has been carried out during annual relief voyages.

The Prince Edward Islands are internationally-important breeding sites for a diversity of seals and seabirds. Most of these species breed during summer months, when marine productivity peaks. Invertebrate and plant activity and growth are also at their highest at this time.<sup>9</sup> Although biological research, including surveys and censuses of biota, does take place at Marion Island during summer (conducted by 'overwintering' personnel spending a year at the island), field teams are then small (normally one to two persons per project), and there are no opportunities to visit Prince Edward Island, 21 km distant.

In December 2001 a summer survey of surface-nesting seabirds was undertaken at both islands (and of fur seals (*Arctocephalus* spp.) and alien plants at Prince Edward Island), resulting in a suite of publications that has been used to support South Africa's national and international obligations to manage the islands effectively.<sup>10,11</sup> In order to assess changes (especially in population sizes) that may have occurred since then, a second summer survey of fur seals and seabirds breeding on the Prince Edward Islands took place in December 2008, with the addition of research on macro-invertebrates and plants at Prince Edward Island and of an oceanographic component. This paper describes the main activities undertaken during the survey and highlights its main findings.

### Survey party and itinerary

A 23-person team of biologists and oceanographers under the overall leadership of Robert Crawford, Marine and Coastal Management, formerly of the Department of Environmental Affairs and Tourism, sailed from Cape Town, South Africa aboard the E.R.V. *Africana* on 10 December 2008. Following pre-sailing quarantine procedures and checks, a boot-washing and kit-inspection 'ceremony', in accordance with SANAP (South African National Antarctic Programme) procedures, was undertaken on the southward voyage to ensure that clothing and equipment going ashore would be free of alien propagules.<sup>16</sup> Most equipment and clothing supplied to the Prince Edward Island camping party (see below) was previously unused as an additional quarantine measure.

The islands were reached by dawn on 16 December. A party of 10 persons was landed at Cave Bay, Prince Edward Island, by rigid inflatable boats on that morning, where a tented camp was established. As part of the landing protocol at Prince Edward Island, 11 rodent bait stations (with scented, attractive baits) were set in and around the camp. The ship then proceeded to Marion Island and landed a seven-person party the same day, leaving six persons aboard to undertake oceanographic research in the vicinity of the islands. The two shore parties then divided

into smaller field teams, working from Cave Bay and Hope Stream on Prince Edward Island, and from the weather station and eight coastal field huts on Marion Island. The Marion field teams were joined by several members of the Marion over-wintering team.

The Prince Edward Island party transferred to Marion Island on 22 December. Following departure from the island the rodent bait stations were inspected for chew marks. No signs were found, and no field evidence of rodents was recorded at any point during the expedition to the island, thought to have always been rodent-free.<sup>12</sup> All solid, including human, wastes were removed from the island for disposal in South Africa. To avoid genetic exchange of biota between the two islands, protective clothing and footwear used on Prince Edward Island was left aboard the ship, being replaced by clean items for use on Marion Island.

Field work on Marion Island was completed on 23 December and all shore-based survey personnel rejoined the ship the next day. After a period spent in the lee of the island sheltering from rough seas, the *Africana* sailed for home on 25 December, arriving in Cape Town on 31 December.

### Preliminary achievements

#### Seals

Antarctic *Arctocephalus gazella* and sub-Antarctic *A. tropicalis* fur seals were surveyed at Prince Edward Island. For both species, counts were made of territorial males and the numbers of pups present estimated.<sup>13</sup> One of two inaccessible sections of coastline was surveyed for the first time, with the beaches between McNish Bay and Kent Crater scanned from the cliffs above. A preliminary analysis revealed that the population of Antarctic fur seals has increased since 2001. An extension of breeding colony sites of sub-Antarctic fur seals was observed. Checks of moulting southern elephant seals (*Mirounga leonina*) found very few flipper tags placed on weaned pups at Marion Island in previous years, suggesting limited movement between the two islands.

#### Birds

Preliminary results suggest that, for most species of surface-nesting seabirds (penguins, albatrosses, giant petrels, cormorants and larids), breeding populations have remained relatively stable since 2001<sup>10</sup> at both islands, although for some species better coverage at Prince Edward Island has led to higher figures. The most marked difference was among cliff-nesting sooty albatrosses (*Phoebastria fusca*), where the 2008 census was almost double that of 2001. This species has recently been uplisted to Endangered (<http://www.birdlife.org/datazone/species/index.html?action=SpcHTMDetails.asp&sid=3967&m=0>), and the larger population on Prince Edward Island further emphasises the importance of this island as a breeding site for albatrosses. A single shy/white-capped albatross (*Thalassarche cauta/steadii*) was found incubating on Prince Edward Island among a small colony of grey-headed albatrosses (*T. chrysostoma*). This is the first time this taxon has been found breeding outside Australasia,<sup>14</sup> bringing the number of species of albatrosses known to have bred on the island to six.

Since 1994, there has been a large decrease (20%) in the mass of southern rockhopper penguins (*Eudyptes chrysocome*) returning to Marion Island from their overwintering feeding grounds to breed.<sup>15</sup> This has been associated with decreased breeding success.<sup>16</sup> The 2008 survey confirmed a large and continued decrease in southern rockhopper penguins at Marion Island, from more than 170 000 pairs in 1994/95<sup>17</sup> to fewer than 60 000

pairs at present. Because Rockhopper Penguins at Marion Island do not feed on commercially-exploited prey, their decrease at this and at other localities within Southern Ocean waters is thought to have resulted from environmental change.<sup>18</sup> For example, recent large displacements of prey items off South Africa have led to mismatches in the distributions of breeding localities and prey, and consequentially to substantial population decreases of some seabirds.<sup>19</sup>

The relative abundances of burrowing petrels (families Procellariidae and Oceanitidae) were studied via spot-lighting at night on both islands and by analysing avian remains in sub-Antarctic skua (*Catharacta antarctica*) middens at Prince Edward Island. A preliminary survey of the distribution and density of burrows of the white-chinned petrel (*Procellaria aequinoctialis*) was made at Prince Edward Island, which will enable a first estimate of population size for the entire island. A similar survey is planned for Marion Island.

Time-depth recorders were fitted to nine macaroni penguins (*E. chrysolophus*) brooding small chicks at Funk Bay, Marion Island, of which eight were recovered. Satellite trackers (Platform Transmitter Terminals; PTTs) were fitted to ten (eight brooding) Indian yellow-nosed albatrosses (*T. carteri*) on Prince Edward Island, a species that does not breed at Marion Island. Two PTTs were placed on light-mantled sooty albatrosses (*Phoebastria palpebrata*) on Marion Island, with a further ten left on the island for later placement on this species and on sooty albatrosses.

Ninety-four hours of quantified seabird and marine mammal observations in total were made from the *Africana* during the southern and northern journeys. Almost 2 300 individuals of 45 species of seabirds and 6 species of cetaceans were identified.

#### Invertebrates

The invertebrate research had four major components. First, invertebrate surveys aimed to assess the density of macro-invertebrates as part of an ongoing study to investigate the influence of introduced house mice (*Mus musculus*), present on nearby Marion Island, but not on Prince Edward Island.<sup>12</sup> Second, individuals of all six weevil species<sup>20,21</sup> were collected as part of an assessment of body-size change associated with climate change and predation by mice on Marion and Prince Edward Islands.<sup>12,22</sup> Third, quantitative assessments of springtail densities were made to compare with previous work<sup>23,24</sup> and to determine which alien species were present. Finally, quantitative sampling of caterpillars, vascular plants, peat, and nest material was undertaken around wandering albatross (*Diomedea exulans*) nests for genetic and isotope analyses. Previous work<sup>25</sup> has suggested that wandering albatrosses serve as thermal ecosystem engineers for *Pringleophaga* sp. (Lepidoptera, Tineidae) caterpillars, which are the main bottleneck for nutrient cycling in the terrestrial system. However, mice may also influence the outcome of this interaction, therefore a control on mouse-free Prince Edward Island was essential.

All of the invertebrate research was completed successfully. It is clear that many invasive alien invertebrates common on Marion Island have not yet managed to colonise Prince Edward Island. These include the springtails *Pogonognathellus flavescens* and *Isotomurus* cf. *palustris*, the slug *Deroceras panormitanum*, the wasp *Aphidius matricariae*, the moth *Plutella xylostella*, and the psychodid midge *Psychoda parthenogenetica*.<sup>26</sup>

#### Plants

Research on the vascular plants on Prince Edward Island had three main components. First, we examined changes in the

distributions of alien vascular plant species, and documented new arrivals. Second, we provided information for ground-truthing a proposed vegetation map, with over 150 vegetation quadrats sampled in a spatially explicit manner. Third, we collected samples for analysis of genetic variation and relationships among islands of two indigenous species, the prostrate herb *Acaena magellanica* (Rosaceae) and the cushion plant *Azorella selago* (Apiaceae), and the alien annual meadow grass *Poa annua*. This research was successfully completed.

No new alien vascular plant species were found on Prince Edward Island since the last survey in 2001.<sup>27</sup> However, procumbent pearlwort *Sagina procumbens* (Caryophyllaceae) has expanded its distribution rapidly since then, and now occurs along virtually all coasts, except for the remote and rather barren High Bluff region. Mouse-eared chickweed *Cerastium fontanum* (Caryophyllaceae) also has extended its range on the north-west side of the island, but was not found above the escarpment and on the eastern side. To record landscape-scale changes, fixed-point photographs were taken at previous points photographed casually by past expeditions to the island and at new points expected to show the most future change.

### Oceanography

The Prince Edward Islands are situated in the pathway of the easterly-flowing Antarctic Circumpolar Current. Bordered to the north by the sub-Antarctic Front and to the south by the Antarctic Polar Front, the islands lie within the Polar Frontal Zone which is a transition zone between sub-Antarctic and Antarctic water masses.<sup>28</sup> The islands act as obstacles to the Antarctic Circumpolar Current, causing high levels of spatial and temporal mesoscale variability within the region.<sup>29,30</sup> It has been shown that waters surrounding the islands demonstrate an 'island mass effect' (enhanced phytoplankton production)<sup>31</sup> whereas an alternative explanation is that food (zooplankton) for the islands' predators (specifically birds) is brought into the vicinity of the islands from upstream regions.<sup>32</sup> It has been suggested that the proximity of the islands to the sub-Antarctic Front affects the availability of zooplankton for predators breeding on the islands,<sup>33</sup> such that the high degree of latitudinal variability of the sub-Antarctic Front may have important implications for food availability for breeding predators.

The main objective was to obtain a summer comparison of the zooplankton community upstream and downstream of the Prince Edward Islands with previous studies conducted during the austral autumn/winter months. Secondary objectives included a comparison of zooplankton abundance and biomass upstream, in the vicinity, and downstream of the islands, to investigate the potential influence of the 'island mass effect'.

Conductivity, temperature and depth casts were conducted at 26 stations (six within the area of the islands and 15 downstream of the islands), with water samples collected at selected depths for oxygen, salinity, chlorophyll-*a* and nutrient analyses. In addition, photosynthesis measurements were conducted, using fast repetition rate fluorometry, on water samples from selected depths within the euphotic zone. The biological component of the oceanographic survey consisted of 21 stations where the mesozooplankton community was sampled using a BONGO net with a 200- $\mu\text{m}$  mesh. Unfortunately, due to unfavourable weather conditions, zooplankton was collected from only nine stations for fatty acid and stable isotope analysis for the determination of trophic links between selected components of plankton and nekton. In addition to net tows, surface seawater was collected with a Crawford bucket at each station for the determination of size-fractionated chlorophyll-*a* concentra-

tions. Surface seawater was collected at three stations for POM (particulate organic matter) analysis. Throughout the survey area, the total zooplankton abundance and biomass was dominated by mesoplankton (200–2000  $\mu\text{m}$ ), comprising mainly copepods. The macrozooplankton community was dominated by chaetognaths, amphipods, euphausiids and salps.

Full results of the 2008 biological survey of the Prince Edward Islands will be published in specialist scientific journals as detailed analyses and interpretation of findings are completed.

The survey was supported financially and logistically by the Branch: Marine and Coastal Management and was conducted with the approval of the Prince Edward Islands Management Committee, Directorate: Antarctica and Islands, then both of the South African Department of Environmental Affairs and Tourism. Funding was also obtained from the National Research Foundation through its South African National Antarctic Programme. Funding for satellite-tracking time was received from the BirdLife International Global Seabird Programme. We thank the following members of the survey team for their willing contributions and support: R.W. Leslie, A. Ouardien, S.L. Petersen, L. Pichegru, M. Slabber, R.J.Q. Tarr, G. Tutt, L. Upfold, J. Visagie, L.J. Waller, D.A. Whitelaw, P.A. Whittington, C. Wilkie, R. Williamson and P. Woodhead. Captain M. Viljoen and the officers and crew of the F.R.V. *Africana* made the sea journeys in often rough seas bearable, and even enjoyable at times, with their friendship and excellent seamanship and catering. We are grateful to the members of the 65th overwintering team at Marion Island, especially L.J. Clokie, J.B. Harding, M.G.W. Jones, P. Kritzing, M.H. Mashau, G.T.W. McClelland, E.M. Mertz and A.M. Treasure, for their help with surveys and hospitality. P.W. Froneman commented on a draft text.

Received 19 February. Accepted 1 July 2009.

- De Villiers M.S. and Cooper J. (2008). Conservation and management. In *The Prince Edward Islands: Land-sea Interactions in a Changing Ecosystem*, eds S.L. Chown and P.W. Froneman, pp. 301–330. African SunMedia, Stellenbosch.
- Crawford R.J.M., Cooper J., du Toit M., Greyling M.D., Hanise B., Holness C.L., Keith D.G., Nel J.L., Petersen S.L., Spencer K., Tshingana D. and Wolfaardt A.C. (2003). Population and breeding of the gentoo penguin *Pygoscelis papua* at Marion Island, 1994/95–2002/03. *Afr. J. Mar. Sci.* **25**, 463–474.
- Crawford R.J.M., Cooper J. and Dyer B.M. (2003). Population of the macaroni penguin *Eudyptes chrysolophus* at Marion Island, 1994/95–2002/03, with observations on breeding and diet. *Afr. J. Mar. Sci.* **25**, 475–486.
- Prince Edward Islands Management Plan Working Group (1996). *Prince Edward Islands Management Plan*. Department of Environmental Affairs and Tourism, Pretoria.
- Department of Environmental Affairs and Tourism (2006). *Prince Edward Islands Environmental Management Plan. Version 0.1*. Prepared for the Department of Environmental Affairs and Tourism, Directorate Antarctica and Islands, by the DST-NRF Centre of Excellence for Invasion Biology, Stellenbosch University, South Africa.
- Davies S.J., Chown S.L. and Joubert L.S. (2007). Renewed management system and provisions for South Africa's sub-Antarctic islands. *Pap. Proc. Roy. Soc. Tasmania* **141**, 115–120.
- Lombard A.T., Reyers B., Schonegevel L.Y., Cooper J., Smith-Adao L.B., Nel D.C., Froneman P.W., Anson I.J., Bester M.N., Tosh C.A., Strauss T., Akkers T., Gon O., Leslie R.W. and Chown S.L. (2007). Conserving pattern and process in the Southern Ocean: designing a Marine Protected Area for the Prince Edward Islands. *Antarct. Sci.* **19**, 39–54.
- Nel D.C. and Ouardien A. (eds) (2008). Towards the development of a Marine Protected Area at the Prince Edward Islands. *WWF South Africa Report Series – 2008/Marine/001*.
- Chown S.L. and Froneman P.W. (eds) (2008). *The Prince Edward Islands: Land-Sea Interactions in a Changing Ecosystem*. African SunMedia, Stellenbosch.
- Cooper J. (ed.) (2003). Seabirds and seals at the Prince Edward Islands. *Afr. J. Mar. Sci.* **25**, 415–562.
- Crawford R.J.M., Underhill L.G., Bester M.N., Cooper J., Greyling M.D., Meyer M.A., Petersen S.L., Ryan P.G. and Wilkinson W. (2003). The Prince Edward Islands Millennium Expedition. *S. Afr. J. Sci.* **99**, 407–409.
- Chown S.L. and Smith V.R. (1993). Climate change and the short-term impact of feral house mice at the sub-Antarctic Prince Edward Islands. *Oecologia* **96**, 508–516.
- Bester M.N., Ryan P.G. and Dyer B.M. (2003). Population numbers of fur seals at Prince Edward Island, Southern Ocean. *Afr. J. Mar. Sci.* **25**, 549–554.
- Brooke M. (2004). *Albatrosses and Petrels around the World*. Oxford University Press, Oxford.
- Crawford R.J.M., Dyer B.M., Cooper J. and Underhill L.G. (2006). Breeding numbers and success of *Eudyptes* penguins at Marion Island, and the influence of mass and time of arrival of adults. *CCAMLR Sci.* **13**, 175–190.



16. Crawford R.J.M., Makhado A.B., Upfold L. and Dyer B.M. (2008). Mass on arrival of rockhopper penguins at Marion Island correlated with breeding success. *Afr. J. Mar. Sci.* **30**, 185–188.
17. Crawford R.J.M., Cooper J., Dyer B.M., Greyling M.D., Klages N.T.W., Nel D.C., Nel J.L., Petersen S.L. and Wolfaardt A.C. (2003). Decrease in numbers of the eastern rockhopper penguin *Eudyptes chrysocome filholi* at Marion Island, 1994/95–2002/03. *Afr. J. Mar. Sci.* **25**, 487–498.
18. Hilton G.M., Thompson D.R., Sagar P.M., Cuthbert R.J., Chereil Y. and Bury S.J. (2006). A stable isotopic investigation into the causes of decline in a sub-Antarctic predator, the rockhopper penguin *Eudyptes chrysocome*. *Glob. Change Biol.* **12**, 611–625.
19. Crawford R.J.M., Sabarros P.S., Fairweather T., Underhill L.G. and Wolfaardt A.C. (2008). Implications for seabirds off South Africa of a long-term change in the distribution of sardine. *Afr. J. Mar. Sci.* **30**, 177–184.
20. Chown S.L. (1992). A preliminary analysis of sub-Antarctic weevil assemblages: local and regional patterns. *J. Biogeogr.* **19**, 87–98.
21. Grobler G.C., Janse van Rensburg L., Bastos A.D.S., Chimimba C.T. and Chown S.L. (2006). Molecular and morphometric assessment of *Ectemnorhinus* weevil species (Coleoptera, Curculionidae, Brachycerinae) from the sub-Antarctic Prince Edward Islands. *J. Zool. Syst. Evol. Res.* **44**, 200–211.
22. Chown S.L. and Klok C.J. (2003). Altitudinal body size clines: latitudinal effects associated with changing seasonality. *Ecography* **26**, 445–455.
23. Gabriel A.G.A., Chown S.L., Barendse J., Marshall D.J., Mercer R.D., Pugh P.J.A. and Smith V.R. (2001). Biological invasions on Southern Ocean islands: the Collembola of Marion Island as a test of generalities. *Ecography* **24**, 421–430.
24. Hugo E.A., Chown S.L. and McGeoch M.A. (2006). The microarthropods of Prince Edward Island: a quantitative assessment. *Polar Biol.* **30**, 109–119.
25. Sinclair B.J. and Chown S.L. (2006). Caterpillars benefit from thermal ecosystem engineering by wandering albatrosses on sub-Antarctic Marion Island. *Biol. Lett.* **2**, 51–54.
26. Chown S.L. and Marshall D.J. (2008). Appendix VII. Terrestrial invertebrates of the Prince Edward Islands. In *The Prince Edward Islands: Land–Sea Interactions in a Changing Ecosystem*, eds S.L. Chown and P.W. Froneman, pp. 400–409. African SunMedia, Stellenbosch.
27. Ryan P.G., Smith V.R. and Gremmen N.J.M. (2003). The distribution and spread of alien vascular plants on Prince Edward Island. *Afr. J. Mar. Sci.* **25**, 555–562.
28. Froneman P.W. and Ansorge I.J. (1998). The third Marion Island Oceanographic Study (MIOS III) conducted in April and May 1998. *S. Afr. J. Sci.* **94**, 937–939.
29. Ansorge I.J., Froneman P.W., Pakhomov E.A., Lutjeharms J.R.E., Perissinotto R. and van Ballegooyen R.C. (1999). Physical-biological coupling in the waters surrounding the Prince Edward Islands (Southern Ocean). *Polar Biol.* **21**, 135–145.
30. Pakhomov E.A. and Froneman P.W. (1999). The Prince Edward Islands pelagic ecosystem, Southern Ocean: a review of achievements. *J. Mar. Syst.* **18**, 355–367.
31. Allanson B.R., Boden B.P. and Parker L. (1985). A contribution to the oceanology of the Prince Edward Islands. In *Antarctic Nutrient Cycles and Food Webs*, eds W.R. Siegfried, P.R. Condy and R.M. Laws, pp. 38–45. Springer, Berlin, Heidelberg, New York.
32. Perissinotto R. (1989). The structure and diurnal variations of the zooplankton of the Prince Edward Islands: implications for the biomass build up of higher trophic levels. *Polar Biol.* **9**, 505–510.
33. Froneman P.W. and Pakhomov E.A. (1998). Biogeographic study of the plankton communities of the Prince Edward Islands. *J. Plankton Res.* **20**, 653–669.