

Fynbos Proteaceae as model organisms for biodiversity research and conservation

Authors:

Frank M. Schurr^{1,2}
Karen J. Esler³
Jasper A. Slingsby⁴
Nicky Allsopp⁴

Affiliations:

¹Plant Ecology and Nature Conservation, Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

²Institut des Sciences de l'Évolution, UMR-CNRS 5554, Université Montpellier II, Montpellier, France

³Department of Conservation Ecology and Entomology, and Centre for Invasion Biology, Stellenbosch University, Stellenbosch, South Africa

⁴South African Environmental Observation Network Fynbos Node, Newlands, Cape Town, South Africa

Correspondence to:

Karen Esler

Email:

kje@sun.ac.za

Postal address:

Private Bag X1, Matieland
7602, South Africa

How to cite this article:

Schurr FM, Esler KJ, Slingsby JA, Allsopp N. Fynbos Proteaceae as model organisms for biodiversity research and conservation. *S Afr J Sci.* 2012;108(11/12), Art. #1446, 4 pages.
<http://dx.doi.org/10.4102/sajs.v108i11/12.1446>

© 2012. The Authors.
Licensee: AOSIS
OpenJournals. This work
is licensed under the
Creative Commons
Attribution License.

Woody plants of the Proteaceae family are a symbol of fynbos. Of the approximately 360 southern African species, over 330 are restricted to the Fynbos biome¹ and form an important part of this biome's exceptional plant diversity.² Proteaceae dominate the overstorey of fynbos vegetation, play a key role for water, carbon and nutrient cycling, and provide resources for many species of pollinators and herbivores.^{1,3} Moreover, Proteaceae are responsible for the bulk of the economic value generated by the fynbos wildflower industry⁴ and serve as flagship species for conservation.

The key role of Proteaceae for the functioning, conservation and economic use of fynbos has led scientists, conservation managers and volunteers to collect a wealth of information on the geographical distribution, ecology and evolutionary history of this group. The foundation for this knowledge was laid by intense research on the population biology of Proteaceae conducted in the 1980s.^{3,5} This research concentrated on 'serotinous' species (36% of fynbos Proteaceae, including most overstorey species, Figure 1a) that store their seeds in fire-safe woody cones and therefore form 'canopy seed banks' but no persistent soil seed banks. Fire triggers the release of seeds from the cones, limiting dispersal and seedling establishment to a short period post-fire. Moreover, 86% of fynbos Proteaceae are 'nonsprouters' (Figure 1b), meaning that fire kills adult plants. In serotinous and nonsprouting Proteaceae, long-term birth, death and population growth rates can be quantified much more easily than for most other plants. Consequently, this group is ideal for studying basic ecological and evolutionary questions. The excellent quantitative understanding of their demography has direct implications for conservation: CapeNature and SANParks use serotinous nonsprouting Proteaceae as indicators to assess the impacts of fire on fynbos biodiversity.

The publications resulting from the Fynbos Biome Project (a National Programme for Ecosystem Research initiated in 1977 by the Council for Scientific and Industrial Research) led to the number of publications on fynbos Proteaceae peaking in 1991 (Figure 2). Since then, interest in Proteaceae has been revived by the Protea Atlas Project, resulting in a second peak in publications on this family in 2010–2011 (Figure 2; in both cases note the long lag times). This remarkable citizen science project involved more than one thousand volunteers assembling about 250 000 estimates of the local abundance of Proteaceae species. This data set paints a detailed picture of the geographical ranges and abundance variation of the fynbos Proteaceae. The Protea Atlas Database has thus been termed 'the largest and highest quality of its kind in the world for studying biodiversity'⁶. It played a crucial role in the development of species distribution models,^{7,8,9} which relate data on species' geographical distributions to environmental variables and have become the predominant method for climate change biodiversity assessments.¹⁰ The database also served as an important source of information for systematic conservation planning.¹¹ Recently, this demographic and biogeographical knowledge has been complemented by molecular data which elucidate the evolutionary processes that created the extraordinary interspecific and intraspecific diversity of Proteaceae.^{12,13,14} As a result of this comprehensive ecological, biogeographical and evolutionary knowledge, the fynbos Proteaceae are now recognised worldwide as model organisms for biodiversity research and conservation.

Recent work building on this exceptional knowledge was presented at this year's Fynbos Forum held from 16 July to 19 July 2012 at Cape St. Francis (South Africa). Researchers, managers, landowners and other stakeholders convened for the special symposium 'Proteaceae: research and management priorities in a changing world'. Presentations at the meeting followed three general themes: the demography of Proteaceae under environmental change; drivers of the group's extraordinary diversity; and interactions between Proteaceae and other organisms.

Proteaceae demography has drawn renewed interest because of the need to understand the dynamics of species and communities under global change. Desika Moodley (Stellenbosch University) measured cone production and seedling recruitment in different populations of *Hakea salicifolia* to understand under which conditions this Australian proteoid becomes naturalised in fynbos. With the aim of informing improved forecasts of native Proteaceae range dynamics




FIGURE 1: Functional diversity of fynbos Proteaceae and representation of the family in scientific publications. The bars give the proportion of study species and fynbos Proteaceae with different (a) seed storage strategies, (b) fire survival strategies and (c) heights.




FIGURE 2: Temporal dynamics of publications on South African Proteaceae. Publication boosts coincided with the end of major research projects (Web of Science search on 13 June 2012 for articles in the fields of Plant Sciences, Ecology, Evolutionary Biology, Biodiversity Conservation or Environmental Sciences that mention Proteaceae or any of the genera occurring in South Africa; relevance of the retrieved publications was then assessed individually). The inset shows the king protea (*Protea cynaroides*), South Africa's national flower (photo[©]: F. Schurr).

in response to changes in climate, land use and fire,^{15,16} John Silander (University of Connecticut, USA), Frank Schurr and Jörn Pögel (Universities of Montpellier, France and Potsdam, Germany) presented studies of large-scale variation in growth, reproduction and mortality across the geographical ranges of several species. Anthony Rebelo (South African National Biodiversity Institute) presented experimental tests of climate change impacts on the germination and seedling emergence of several Proteaceae species. Germination was reduced at higher temperatures, but some seeds still germinated at temperatures 3.5 °C higher than at present. Seedlings differed strongly in their tolerance of extended drought although the causes are poorly understood.¹⁷ While climate change effects on Proteaceae demography are still under-researched, certain populations appear threatened by increases in fire frequency observed in much of the Fynbos biome.¹⁸ Three demographic studies performed by Annelise Schutte-Vlok (CapeNature), Tineke Kraaij (SANParks) and Katharina Mayer (Potsdam, Germany), independently concluded that contemporary fire intervals are frequently too short for slow-maturing nonsprouting species to set seed. Small-scale demographic variation is also the focus of ongoing research presented by Henning Nottebrock (Potsdam, Germany) and Baptiste Schmid (Biodiversity and Climate Research Centre, Germany). They mapped all shrub individuals in 27 *Protea*-dominated communities at high spatial resolution, quantifying how plant traits determine the strength and spatial scale of plant–plant and plant–pollinator interactions and how these interactions affect the reproduction of individual plants.

Potentially, such trait-based demographic studies will not only improve our understanding of existing communities, but will also help to predict the dynamics of novel communities that will arise when global change reshuffles the geographical distribution of species.¹⁹ Proteaceae are well suited for such trait-based studies as they show remarkable variation in plant size, leaf form, ability to survive fires and many other functional traits. This variation is the reason Linnaeus named the family after the Greek god Proteus, who could take any shape.

The causes of this extraordinary functional diversity were a second cross-cutting theme at the meeting. Several studies focused on leaf traits (e.g. leaf size, hairiness and specific leaf area), which show adaptive differentiation among populations^{20,21} and may determine the vulnerability of species to climate change.²² Using transplant experiments, Jane Carlson and Kent Holsinger (Connecticut, USA) showed that populations from extremely cold and dry environments have the least plastic leaf traits, suggesting that these extreme populations have the lowest potential for rapid, plastic responses to environmental change. Ongoing genomic analyses should elucidate the causes of such intraspecific differentiation. Studying larger timescales, Byron Lamont (Curtin University, Australia) and Jeanne Tonnabel (Montpellier) respectively



investigated the evolution of serotiny in the genera *Protea* and *Leucadendron*. These studies independently suggest that serotiny is ancestral in both genera and that fire shaped their evolution at least 28 million years ago. Family-wide macroevolutionary studies even suggest that the Proteaceae developed fire adaptations as early as 88 million years ago.²³ Lamont also discussed a recently launched project that aims to construct a species-level phylogeny of all Proteoideae, the subfamily containing most native African Proteaceae. Such phylogenies are important in understanding how diversity in the Proteaceae arose and was maintained in order to potentially inform predictions of future biodiversity dynamics.²⁴

A third line of research examined pollination, seed dispersal and seed predation interactions. Sandy-Lynn Steenhuisen (University of Cape Town) showed that rodent-pollinated *Protea* species evolved repeatedly from bird-pollinated ancestors and that these pollinator shifts are associated with the appearance of sour-milk floral scents. Such pollinator shifts can even occur within species: Christopher Johnson (Stellenbosch University) showed that the common form of *Leucospermum tottum* can be pollinated by both birds and insects, whereas a narrowly distributed variety relies on birds. Using experiments and pollinator observations in mapped communities, Baptiste Schmid quantified how strongly plants depend on bird pollination and compete for bird visits. Anina Heystek (Stellenbosch University) suggested that pollinator-mediated competition even shapes large-scale plant distributions: analysing *Protea* Atlas data, she showed that Proteaceae with the same pollination syndrome co-occur less frequently than expected by chance. Finally, Ursina Rusch (Stellenbosch University) studied rodents which cache and eat nut-like Proteaceae seeds, thereby acting as both seed disperser and seed predator. The Cape spiny mouse (*Acomys subspinosus*) caches single seeds, thereby hiding them from the seed-predating striped mouse (*Rhabdomys pumilio*) which poorly detects single seeds.

The scientific presentations were followed by a workshop attended by 37 people from a range of research and management backgrounds. When asked: 'What are the most important questions from a management perspective?', participants primarily mentioned the topics fire (38%), conservation planning (23%), indicators and monitoring (13%) and climate change (12%). Researchers tended to ask questions which were basic in nature (64% basic vs 36% applied, $N = 22$) whilst the opposite was true for managers (28% basic vs 72% applied, $N = 15$). So even within the bounds of the set question, researchers and managers framed questions for applied research differently. This difference emphasises the importance of encouraging opportunities for researchers and managers to develop research priorities in collaboration. Yet it was encouraging that ongoing research initiatives aim to address many of the applied questions asked. For example, to assess biodiversity impacts of fire management, wild flower harvesting and habitat

fragmentation, one needs demographic knowledge (such as how reproduction and mortality depend on fire intervals, which populations are seed-limited and how long isolated populations can persist).

It could be argued that fynbos research has focused too strongly on the Proteaceae, neglecting most of the biome's plant diversity. Even within the Proteaceae, non-serotinous and small-statured species are understudied (Figures 1b and 1c), and the full research potential provided by the functional variation of this iconic group is yet to be realised. Nevertheless, concentrating on a single taxon enables the quantification of regional variation and should yield nuanced insights for conservation across the biome. Although broadening the scope to provide better coverage of other fynbos families will be invaluable, we expect that Proteaceae will continue to play a central role for biodiversity research in the fynbos and beyond.

Acknowledgements

The special symposium and workshop on Proteaceae were held in celebration of the German–South African Year of Science, and were funded by a grant of the German Federal Ministry of Education and Research, the South African Department of Science and Technology and the National Research Foundation to K.J.E., N.A. and F.M.S. Additionally, F.M.S. acknowledges support from the German Research Foundation (SCHU 2259/3-1 and SCHU 2259/5-1). Kent Holsinger, Tineke Kraaij, Byron Lamont, Jörn Pagel and the rest of the workshop participants are thanked for their valuable inputs.

References

1. Rebelo AG. *Proteas: A field guide to the proteas of southern Africa*. Vlaeberg: Fernwood Press; 2001.
2. Goldblatt P, Manning JC. Plant diversity of the Cape region of southern Africa. *Ann Missouri Bot Gard*. 2002;89:281–302. <http://dx.doi.org/10.2307/3298566>
3. Cowling RM, editor. *The ecology of fynbos: Nutrients, fire and diversity*. Cape Town: Oxford University Press; 1992.
4. Turpie JK, Heydenrych BJ, Lamberth SJ. Economic value of terrestrial and marine biodiversity in the Cape Floristic Region: Implications for defining effective and socially optimal conservation strategies. *Biol Conserv*. 2003;112:233–251. [http://dx.doi.org/10.1016/S0006-3207\(02\)00398-1](http://dx.doi.org/10.1016/S0006-3207(02)00398-1)
5. Bond WJ, Van Wilgen BW. *Fire and plants*. London: Chapman & Hall; 1996. <http://dx.doi.org/10.1007/978-94-009-1499-5>
6. Gelfand AE, Schmidt AM, Wu S, et al. Modelling species diversity through species level hierarchical modelling. *Journal of the Royal Statistical Society Series C*. 2005;54:1–20. <http://dx.doi.org/10.1111/j.1467-9876.2005.00466.x>
7. Midgley GF, Hannah L, Millar D, et al. Assessing the vulnerability of species richness to anthropogenic climate change in a biodiversity hotspot. *Global Ecol Biogeogr*. 2002;11:445–451. <http://dx.doi.org/10.1046/j.1466-822X.2002.00307.x>
8. Thomas CD, Cameron A, Green RE, et al. Extinction risk from climate change. *Nature*. 2004;427:145–148. <http://dx.doi.org/10.1038/nature02121>
9. Latimer AM, Wu SS, Gelfand AE, et al. Building statistical models to analyze species distributions. *Ecol Appl*. 2006;16:33–50. <http://dx.doi.org/10.1890/04-0609>
10. Dawson TP, Jackson ST, House JL, et al. Beyond predictions: Biodiversity conservation in a changing climate. *Science*. 2011;332:53–58. <http://dx.doi.org/10.1126/science.1200303>
11. Cowling RM, Pressey RL, Rouget M, et al. A conservation plan for a global biodiversity hotspot – the Cape Floristic Region, South Africa. *Biol Conserv*. 2003;112:191–216. [http://dx.doi.org/10.1016/S0006-3207\(02\)00425-1](http://dx.doi.org/10.1016/S0006-3207(02)00425-1)
12. Barker NP, Vanderpoorten A, Morton CM, et al. Phylogeny, biogeography and the evolution of life-history traits in *Leucadendron* (Proteaceae). *Mol Phylogenet Evol*. 2004;33:845–860. <http://dx.doi.org/10.1016/j.ympev.2004.07.007>



13. Valente LM, Reeves G, Schnitzler J, et al. Diversification of the African genus *Protea* (Proteaceae) in the Cape biodiversity hotspot and beyond: Equal rates in different biomes. *Evolution*. 2010;64:745–759. <http://dx.doi.org/10.1111/j.1558-5646.2009.00856.x>
14. Prunier R, Holsinger KE, Carlson JE. The effect of historical legacy on adaptation: Do closely related species respond to the environment in the same way? *J Evol Biol*. 2012;25:1636–1649. <http://dx.doi.org/10.1111/j.1420-9101.2012.02548.x>
15. Pagel J, Schurr FM. Forecasting species ranges by statistical estimation of ecological niches and spatial population dynamics. *Global Ecol Biogeogr*. 2012;21:293–304. <http://dx.doi.org/10.1111/j.1466-8238.2011.00663.x>
16. Schurr FM, Pagel J, Sarmiento Cabral J, et al. How to understand species' niches and range dynamics: A demographic research agenda for biogeography. *J Biogeogr*. 2012; doi:10.1111/j.1365-2699.2012.02737.x. <http://dx.doi.org/10.1111/j.1365-2699.2012.02737.x>
17. Mustart PJ, Rebelo AG, Juritz J, et al. Wide variation in post-emergence desiccation tolerance of seedlings of fynbos proteoid shrubs. *S Afr J Bot*. 2012;80:110–117. <http://dx.doi.org/10.1016/j.sajb.2012.03.009>
18. Wilson AM, Latimer AM, Silander JA, et al. A hierarchical Bayesian model of wildfire in a Mediterranean biodiversity hotspot: Implications of weather variability and global circulation. *Ecol Model*. 2010;221:106–112. <http://dx.doi.org/10.1016/j.ecolmodel.2009.09.016>
19. Lavergne S, Mouquet N, Thuiller W, et al. Biodiversity and climate change: Integrating evolutionary and ecological responses of species and communities. *Annu Rev Ecol Evol S*. 2010;41:321–350. <http://dx.doi.org/10.1146/annurev-ecolsys-102209-144628>
20. Carlson JE, Holsinger KE, Prunier R. Plant responses to climate in the Cape Floristic Region of South Africa: Evidence for adaptive differentiation in the Proteaceae. *Evolution*. 2011;65:108–124. <http://dx.doi.org/10.1111/j.1558-5646.2010.01131.x>
21. Prunier R, Holsinger KE. Was it an explosion? Using population genetics to explore the dynamics of a recent radiation within *Protea* (Proteaceae L.). *Mol Ecol*. 2010;19:3968–3980. <http://dx.doi.org/10.1111/j.1365-294X.2010.04779.x>
22. Skelton RP, Midgley JJ, Nyaga JM, et al. Is leaf pubescence of Cape Proteaceae a xeromorphic or radiation-protective trait? *Aust J Bot*. 2012;60:104–113. <http://dx.doi.org/10.1071/BT11231>
23. Lamont BB, He T. Fire-adapted Gondwanan angiosperm floras evolved in the Cretaceous. *BMC Evol Biol*. In press; 2012.
24. Parallel evolutionary radiations in *Protea* and *Pelargonium* in the greater Cape Floristic Region [homepage on the Internet]. c2012 [updated 2012 Aug 14; cited 2012 Aug 31]. Available from: http://darwin.eeb.uconn.edu/wiki/index.php/Parallel_Evolutionary_radiations_in_Protea_and_Pelargonium_in_the_Greater_Cape_Floristic_Region