

# Antibacterial, antioxidant activities and cytotoxicity of plants against *Propionibacterium acnes*

**AUTHORS:**  
Richa Sharma<sup>1</sup>  
Namrita Lall<sup>1</sup>

**AFFILIATION:**  
<sup>1</sup>Department of Plant Science,  
University of Pretoria, Pretoria,  
South Africa

**CORRESPONDENCE TO:**  
Namrita Lall

**EMAIL:**  
namrita.lall@up.ac.za

**POSTAL ADDRESS:**  
Department of Plant Science,  
University of Pretoria, Private  
Bag X20, Hatfield 0028,  
South Africa

**DATES:**  
**Received:** 16 Sep. 2013  
**Revised:** 03 Jan. 2014  
**Accepted:** 23 Mar. 2014

**KEYWORDS:**  
anti-acne; mouse melanocytes;  
medicinal plants; 2,2-diphenyl-1-  
picrylhydrazal; ethanolic extract

**HOW TO CITE:**  
Sharma R, Lall N. Antibacterial,  
antioxidant activities and  
cytotoxicity of plants against  
*Propionibacterium acnes*.  
S Afr J Sci. 2014;110(11/12),  
Art. #2013-0293, 8 pages.  
[http://dx.doi.org/10.1590/  
sajs.2014/20130293](http://dx.doi.org/10.1590/sajs.2014/20130293)

The use of plants to treat skin ailments has strong support in the current trend of drug discovery. *Propionibacterium acnes*, an anaerobic pathogen, plays an important role in the occurrence of acne. The present study was conducted to evaluate the antimicrobial and antioxidant activities against *P. acnes* and cytotoxic effects of 48 medicinal plants grown in South Africa. The broth dilution and DPPH radical scavenging methods were used to determine antibacterial and antioxidant activities, respectively. Cytotoxicity was determined on mouse melanocytes (B16-F10). The ethanolic bark extract of *Acacia galpinii* Burt Davy. (Leguminosae) exhibited the lowest minimum inhibitory concentration of 62.5 µg/mL. Excellent antioxidant activity was shown by *Aspalathus linearis* (Burm.f.) R.Dahlgren (Leguminosae), *Combretum apiculatum* Sond. (Combretaceae), *Harpephyllum caffrum* Bernh. ex Krauss (Anacardiaceae) and *Sclerocarya birrea* Hochst. (Anacardiaceae), with 50% radical scavenging activity (EC<sub>50</sub>) at concentrations ranging from 1.6 µg/mL to 3.5 µg/mL. *Greyia sutherlandii* Hook. & Harv. (Greyiaceae) also exhibited good antioxidant activity with an EC<sub>50</sub> value of 7.9±0.23 µg/mL. *A. linearis*, *G. sutherlandii* and *S. birrea* showed low toxicity with 50% viability of cells (EC<sub>50</sub>) at concentrations of 125.09±0.71 µg/mL, 107.85±1.53 µg/mL and 92.07±0.09 µg/mL, respectively. The extracts of *A. linearis*, *G. sutherlandii* and *S. birrea* showed good antibacterial and antioxidant activities and low toxicity. Therefore, these plants can be considered as possible anti-acne agents and warrant further investigation.

## Introduction

Acne, one of the most common disorders of the skin, is a polymorphic disease with non-inflammatory (blackhead or whitehead) and inflammatory (papules, pustules, or nodules) aspects and a wide spectrum of severity. Acne can have a significant impact on the psychosocial and physical aspects of life. It affects up to 85% of adolescents to some extent but is less common among infants. Its prevalence has been estimated to be 95–100% in male adolescents and 83–85% in female adolescents.<sup>1-4</sup>

*Propionibacterium acnes*, a Gram-positive anaerobic bacterium, is a normal component of the microbiota of human skin. *P. acnes* causes an increase in the secretion of sebum from sebaceous glands, which is accompanied by the thickening of the epidermis at the outlet to the pilosebaceous follicles. As a result, there is an obstruction to the flow of sebum outwards, and a comedone develops. Colonisation of the follicles with *P. acnes* and the host's inflammatory response play a pivotal role in the development of typical inflammatory papulopustular lesions.<sup>5</sup> In an anaerobic environment, the bacteria secretes nucleases, nурaminidases, hyaluronidases, acid phosphatases, lecithinases and other lipases. As a result of the action of these enzymes, the sebum content changes and reactive oxygen species may be released from the damaged follicular walls. Reactive oxygen species may also be the reason for the progression of inflammation in the pathogenesis of disease.<sup>6</sup>

Conventional drugs commonly used in acne treatment – such as tetracycline, erythromycin, mynocyline and metronidazole – act as antioxidants and antibacterials. Benzoyl peroxide, a topical agent for the treatment of acne, shows the ability to induce an inflammatory reaction mediated by reactive oxygen species in addition to its antibacterial activity.<sup>6</sup> These drugs also have various known side effects. The topical antibiotics can lead to dryness, redness and irritation of the skin, as well as hypopigmentation while oral antibiotics have age restrictions, can cause gastrointestinal disorders and increase the risk of venous thromboembolism.<sup>5</sup>

Herbal medicines are an important part of African tradition and also have very deep roots in the treatment of dermatological ailments. Ethnobotanical studies have documented the use of plants by traditional healers for the treatment of various skin ailments.<sup>7</sup> Different plant parts commonly used as cosmetics or face masks, known as *umemezis*, are widely used in southern Africa for skin problems like inflammation, wounds, burns, eczema and puberty acne.<sup>8</sup>

Because many skin disorders like atopic dermatitis and acne are associated with inflammation and the release of free radicals, which lead to oxidative and cellular damage and bacterial infections such as *P. acnes*, the presence of antioxidant and antimicrobial agents can explain the effectiveness of plants in the treatment of skin infections. In order to develop the therapeutic and drug potential of these plants, it is important to know whether they have any cytotoxic effects. Therefore, ethanol extracts of selected plants were evaluated for their antibacterial and antioxidant activities and cytotoxicity.

Limitations in the usage of some drugs and the prevailing side effects of the various chemically derived compounds have led to the search for alternative herbal agents to treat acne. The aim of this study was to test the effect of selected plant extracts on the pathogenic bacteria *P. acnes*, and to identify which plant extracts could be considered as possible anti-acne agents.

## Methods

### Materials

Tetracycline, vitamin C, *p*-iodonitrotetrazolium salt and 2,2-diphenyl-1-picrylhydrazyl (DPPH) were obtained from Sigma-Aldrich (Johannesburg, South Africa). Nutrient agar and nutrient broth were obtained from Merck

SA (Pty) Ltd (Johannesburg, South Africa). *P. acnes* (ATCC 11827) was purchased from Anatech Company South Africa (Johannesburg, South Africa). The cell culture reagents and the equipment were purchased from Highveld Biological (Johannesburg, South Africa), Labotech (Johannesburg, South Africa) and The Scientific Group (Johannesburg, South Africa). The B16-F10 mouse melanocyte cell line was obtained from Highveld Biological.

### Preparation of plant extracts

Different plant parts (leaves, roots, bark and twigs) were collected from the Botanical Garden of the University of Pretoria (Pretoria, South Africa). The plants were identified by a taxonomist, Professor A.E. (Braam) van Wyk, at the H.G.W.J. Schweickerdt Herbarium of the University of Pretoria. The shade-dried plant material (80 g) was ground with a mechanical grinder, then soaked in 300 mL of ethanol and left on a shaker for 3 days. The plant material was then filtered and the solvent was evaporated under vacuum (Buchi Rotavapor, Labotech, Switzerland) to yield dry extracts. The plants were selected based on their medicinal usage as summarised in Table 1.

### Antibacterial bioassay

The minimal inhibitory concentration (MIC) of the ethanolic extracts of the 48 selected plants was determined by a microdilution assay. This assay was done using the method described by Mapunya et al.<sup>36</sup>, with slight modifications. For this purpose, *P. acnes* (ATCC 11827) was cultured from a Kwik-Stick on nutrient agar and incubated at 37 °C for 72 h under anaerobic conditions. The ethanolic extracts were dissolved in 10% dimethyl sulphoxide (DMSO) to obtain a stock solution of 2 mg/mL. The positive control (tetracycline) was dissolved in sterile distilled water to obtain a stock solution of 0.2 mg/mL. The 96-well plates were prepared by dispensing 100 µL of the nutrient broth into each well; 100 µL of the plant stock samples and positive control were added to the first row of wells in triplicate. Twofold serial dilutions were made in broth over a range to give concentrations of 3.9–500 µg/mL and 0.3–50 µg/mL for the plant extracts and positive control, respectively. The 72-h culture of bacteria was dissolved in nutrient broth and the suspensions were adjusted to 0.5 McFarland standard turbidity at 550 nm. Then 100 µL of this bacterial inoculum with 10<sup>5</sup>–10<sup>6</sup> CFU/mL was added to all the wells. The wells with 2.5% DMSO and bacterial suspension without samples served as the solvent and negative controls, respectively. The plates were then incubated at 37 °C for 72 h under anaerobic conditions. The MIC (defined as the lowest concentration that showed no bacterial growth) was determined by observing the colour change in the wells after the addition of *p*-iodonitrotetrazolium salt.

### Antioxidant assay

The antioxidant activity of selected plant extracts was investigated using the DPPH radical scavenging method as previously described by Du Toit et al.<sup>37</sup>, with slight modifications. DPPH is a free radical, which is stable at room temperature and produces a violet solution in ethanol. When reduced in the presence of an antioxidant molecule, it gives rise to a colourless solution. DPPH was dissolved in ethanol to obtain a solution of 0.04% w/v.

The selected plant samples and the positive control (vitamin C) stock solutions (2 mg/mL) were serially diluted to final concentrations ranging from 0.78 µg/mL to 100 µg/mL. Ethanol and DPPH without any plant material were used as blanks while plant samples diluted in distilled water were used as controls. DPPH solution (90 µg/mL) was then added to all the wells except for the controls and allowed to react at room temperature. After 30 min, the absorbance values were measured at 515 nm using a Biotek Power-wave XS multiwell reader (A.D.P., Johannesburg, South Africa). The values were converted into the percentage antioxidant activity (AA) using the formula given below. The 50% inhibitory concentration (EC<sub>50</sub>) values were then calculated by linear regression of the plots using GraphPad Prism version 4.

$$AA\% = \left\{ \frac{\text{Abs}_{\text{blank}} - (\text{Abs}_{\text{sample}} - \text{Abs}_{\text{control}})}{\text{Abs}_{\text{blank}}} \right\} * 100$$

### Mouse melanocyte cytotoxicity assay

The cytotoxicity of selected plant extracts was determined following a previously described method.<sup>36</sup> Briefly, mouse melanocyte (B16-F10) cells were plated in complete Roswell Park Memorial Institute medium (10% foetal bovine serum and 1% gentamycin) directly in the wells of a 96-well plate (10<sup>5</sup> cells per well). After an overnight incubation at 37 °C in 5% CO<sub>2</sub> and a humidified atmosphere, extract samples and the positive control (actinomycin D) were added to the cells to give the final concentrations of plant extract and positive control of 3.13–400 µg/mL and 0.03x10<sup>-2</sup>–0.05 µg/mL, respectively. Plates were incubated at 37 °C in 5% CO<sub>2</sub> in a humidified atmosphere for 3 days. The toxicity effects of the extracts on the B16-F10 cells were assayed using the sodium 3'-[1-(phenyl amino-carbonyl)-3,4-tetrazolium]-bis-[4-methoxy-6-nitrobenzene sulphonic acid hydrate (XTT) cytotoxicity assay. Thereafter, 50 µL of XTT reagent (1 mg/mL XTT with 0.383 mg/mL penazine methosulphate) was added to the wells and incubated for 1 h. The optical densities of the wells were measured at 450 nm with background subtraction at 690 nm. Cell survival was assessed by comparison with the controls (medium with DMSO). The EC<sub>50</sub> value, which represents the concentration of plant extract that causes death in 50% of the cells, was analysed using GraphPad Prism version 4.

### Statistical analysis

All the assays were performed in triplicate with three independent studies for each assay. EC<sub>50</sub> values for antioxidant and cytotoxicity tests were derived from a non-linear regression model (curve fit) based on a sigmoidal dose response curve (variable) and computed using GraphPad Prism version 4 (GraphPad, San Diego, CA, USA).

## Results

### Antibacterial activity of ethanolic extracts

The antibacterial activity of the selected plants against *P. acnes* is summarised in Table 2. After the addition of *p*-iodonitrotetrazolium salt, the MIC value of the positive drug control (tetracycline) was determined to be 3.1 µg/mL. Most of the plants exhibited antibacterial activity at MIC values ranging from 62.5 µg/mL to 500 µg/mL. The ethanolic bark extract of *Acacia galpinii* Burt Davy. (Leguminosae) exhibited the lowest MIC value of 62.5 µg/mL. Some of the plant extracts – *Aspalathus linearis* (Burm.f.) R.Dahlgren (Leguminosae), *Combretum apiculatum* Sond. (Combretaceae), *Combretum molle* Engl. & Diels (Combretaceae), *Galenia africana* L. (Aizoaceae), *Greyia sutherlandii* Hook. & Harv. (Greyiaceae), *Harpephyllum caffrum* Bernh. ex Krauss (Anacardiaceae), *Ranunculus repens* L. (Ranunculaceae), *Sclerocarya birrea* Hochst. (Anacardiaceae) and *Warburgia salutaris* (G. Bertol.) Chiov. (Cancellaceae) – exhibited MIC values of 125 µg/mL. Another 28 extracts inhibited the growth of bacteria at MIC values ranging from 250 µg/mL to 500 µg/mL, whereas the remaining 10 extracts did not show any antibacterial activity, even at the highest concentration (500 µg/mL) tested. A threshold MIC value of ~100 µg/mL is suggested for rating plant extracts as having significant antimicrobial activity.<sup>38</sup> Therefore, the plant extracts exhibiting MIC values ranging from 62.5 µg/mL to 125 µg/mL were selected for the evaluation of antioxidant activity.

### Antioxidant activity of selected extracts

Vitamin C, a widely used antioxidant compound, was used as the positive control (EC<sub>50</sub> = 1.98 ± 0.005 µg/mL). The plant extracts which demonstrated excellent radical scavenging activity, comparable to vitamin C, were *A. linearis* (EC<sub>50</sub> of 3.5 ± 0.5 µg/mL), *C. apiculatum* (EC<sub>50</sub> of 1.6 ± 0.02 µg/mL), *H. caffrum* (EC<sub>50</sub> of 2.6 ± 0.21 µg/mL) and *S. birrea* (EC<sub>50</sub> of 2.06 ± 0.03 µg/mL) (Figure 1). The plant extracts of *C. molle* and *G. sutherlandii* also showed good antioxidant activity with EC<sub>50</sub> values of 9.83 ± 0.8 µg/mL and 7.9 ± 0.23 µg/mL, respectively (Figure 1). *A. galpinii* and *R. repens* exhibited comparatively higher antioxidant activity with EC<sub>50</sub> values of 16.05 ± 2.25 µg/mL and 24.7 ± 2.05 µg/mL, respectively. The extracts of *G. africana* and *W. salutaris* exhibited the lowest radical scavenging activity with the highest EC<sub>50</sub> values of 90.92 ± 1.2 µg/mL and 111 ± 2.5 µg/mL, respectively.


**Table 1:** Medicinal use of plants selected for present study

Plant Name	Medicinal use
<i>Acacia caffra</i> Willd.	Treatment of blood disorders, infantile abdominal disorders <sup>7</sup>
<i>Acacia galpinii</i> Burt Davy.	As a demulcent <sup>8</sup>
<i>Acacia mellifera</i> Benth.	Treatment of coughs, gastrointestinal ailments, malaria, pneumonia, stomach aches, sterility, skin diseases <sup>9</sup>
<i>Aloe arborescens</i> Mill.	Used in cosmetics; treatment of X-ray burns, stomach aches <sup>7</sup>
<i>Aloe barbadensis</i> Mill.	As an antioxidant; used in cosmetic application, wound healing <sup>7,10,11</sup>
<i>Aloe ferox</i> Mill.	Treatment of ophthalmia, venereal sores <sup>7</sup>
<i>Aloe sessiliflora</i> Pole-Evans.	Believed to promote menstruation; as enemas <sup>7,12</sup>
<i>Anchusa capensis</i> Thunb.	Used as a mutagenic and neurotoxin, and in traditional phytomedicine <sup>13</sup>
<i>Annona senegalensis</i> Pers.	Treatment of dermatological diseases and ophthalmic disorders <sup>14</sup>
<i>Arbutus unedo</i> L.	As anti-diarrhoeal, astringent, antioxidant, urinary antiseptic, depurative; treatment of diabetes and hypertension <sup>15-18</sup>
<i>Aspalathus linearis</i> (Burm.f.) R.Dahlgren	Used for alleviation of infantile colic, allergies, asthma, dermatological problems <sup>12</sup>
<i>Barleria albostellata</i> C. B. Clarke	As antibacterial, antifungal, anti-inflammatory, antioxidant and for acne inhibition <sup>19</sup>
<i>Barleria repens</i> Nees	As antibacterial, antifungal <sup>19</sup>
<i>Broussonetia papyrifera</i> (L.) Vent.	Treatment of stomach pains, ill-defined abdominal pains <sup>20</sup>
<i>Buxus macowanii</i> Oliv.	Treatment of gout, malaria, rheumatism, skin disorders <sup>13</sup>
<i>Carpobrotus edulis</i> (L.) Bolus	Treatment of infections of the mouth and throat, eczema, wounds and burns <sup>12</sup>
<i>Ceratonia siliqua</i> L.	As anti-diarrhoeal, antitussive, diuretic; treatment of warts <sup>21</sup>
<i>Combretum apiculatum</i> Sond.	Treatment of conjunctivitis, stomach disorders <sup>7</sup>
<i>Combretum molle</i> Engl. & Diels	As a anthelmintic; treatment of coughs, fever, stomach ailments, wounds <sup>7</sup>
<i>Cotyledon orbiculata</i> L.	Treatment of earache, toothache, epilepsy, boils and inflammation <sup>7,12</sup>
<i>Cryptocarya woodii</i> Engl.	Treatment of diarrhoea <sup>22</sup>
<i>Dahlia imperialis</i> Roehl	Treatment of skin ailments like rashes, grazes, infected scratches <sup>23</sup>
<i>Datura stramonium</i> L.	Treatment of abscesses and wounds; relieves asthma and reduces pain; remedy for boils <sup>7,12</sup>
<i>Dichrostachys cinerea</i> (L.) Wight & Arn.	Treatment of body pains, elephantiasis, sores and skin ailments, toothache <sup>7,12</sup>
<i>Diospyros lycioides</i> Desf.	Chewed and used as a toothbrush, and to ease body pains <sup>7</sup>
<i>Dodonaea viscosa</i> Jacq.	As antipruritic in skin rashes and fungal skin diseases <sup>12</sup>
<i>Erythrophleum lasianthum</i> Corbishley.	Treatment of fever, general body pains, headaches, intestinal spasms, migraines <sup>7,12</sup>
<i>Euclea divinorum</i> Hiern.	As chewing sticks for toothache, headaches; as a purgative; bark infusion is used to enhance appetite <sup>7</sup>
<i>Euclea natalensis</i> A.DC.	Treatment of bronchitis, chronic asthmas, pleurisy, toothache, urinary tract infections <sup>7</sup>
<i>Galenia africana</i> L.	Treatment of asthma, coughs, skin diseases, eye inflammation, venereal sores, wounds <sup>24</sup>
<i>Gomphocarpus fruticosus</i> R.Br.	Treatment of headache and tuberculosis and to relieve stomach pain and general aches in the body <sup>7,12</sup>
<i>Greyia flanaganii</i> Bolus.	To ward off sickness <sup>25</sup>
<i>Greyia sutherlandii</i> Hook. & Harv.	As emetics for biliousness <sup>7</sup>
<i>Harpephyllum caffrum</i> Bernh. ex Krauss	As blood purifiers, for facial saunas, skin washes; to treat acne and eczema <sup>7,12</sup>
<i>Heteropyxis natalensis</i> Harv.	Treatment of colds, bleeding gums, nosebleeds; as a vermifuge <sup>7,12</sup>
<i>Hyaenanche globosa</i> Lamb.	Treatment against vermins <sup>13</sup>
<i>Knowltonia vesicatoria</i> Sims.	Treatment of headaches, toothaches, skin blisters <sup>12</sup>
<i>Magnolia grandiflora</i> L.	Treatment of abdominal discomfort, blood pressure, dyspnoea, epilepsy, heart disturbances, infertility, muscle spasm <sup>26,27</sup>
<i>Myrsine africana</i> L.	As anthelmintics, blood purifier <sup>28</sup>
<i>Parinari curatellifolia</i> Planch. ex Benth.	Treatment of ailments of the eye or ear, pneumonia <sup>29,30</sup>
<i>Ranunculus repens</i> L.	Treatment of muscular aches, rheumatic pains, sores <sup>31</sup>
<i>Rhus lancea</i> L.f.	As antibacterial and antifungal <sup>32</sup>
<i>Sclerocarya birrea</i> Hochst.	Treatment of diarrhoea, inflammation, skin ailments, stomach ailments, malaria, ulcers <sup>7,12</sup>
<i>Sideroxylon inerme</i> L.	As skin lightener; to treat fevers; to treat gall sickness in stock <sup>7,12,33</sup>
<i>Symphytum officinale</i> L.	Treatment of arthritis, bruises, insect bites, inflamed bunions, wounds, skin conditions, nosebleeds, sunburn, rheumatism <sup>34</sup>
<i>Warburgia salutaris</i> (G. Bertol.) Chiov.	Treatment of influenza, rheumatism, malaria, venereal diseases, headaches, toothaches, dermatological disorders, gastric ulcers <sup>7,12</sup>
<i>Zanthoxylum capense</i> Harv.	Treatment of colic, stomach aches, toothaches, fever, epilepsy <sup>7,12,35</sup>

**Table 2:** Minimum inhibitory concentrations (MICs) for antibacterial activity of extracts against *Propionibacterium acnes* determined by microdilution assay

Plant name	Common name	Family	Voucher no.	Part used	MIC µg/mL
<i>Acacia caffra</i>	Cat thorn	Leguminosae	PRU 90700	Leaves	250
<i>Acacia galpinii</i>	Monkey thorn	Leguminosae	PRU 16209	Bark	62.5
<i>Acacia mellifera</i>	Blackthorn	Leguminosae	PRU 078373	Leaves	250
<i>Aloe arborescens</i>	Krantz aloe	Aloaceae	MN 5	Leaves	500
<i>Aloe barbadensis</i>	Aloe vera	Aloaceae	PRU 118947	Leaves	Na <sup>†</sup>
<i>Aloe ferox</i>	Red aloe	Aloaceae	PRU 110308	Leaves	Na <sup>†</sup>
<i>Aloe sessiliflora</i>	Lebombo aloe	Aloaceae	PRU 118948	Leaves	Na <sup>†</sup>
<i>Anchusa capensis</i>	Cape forget-me-not	Boraginaceae	Not available	Leaves	Na <sup>†</sup>
<i>Annona senegalensis</i>	White custard apple	Annonaceae	PRU 074974	Bark	250
<i>Arbutus unedo</i>	Strawberry tree	Ericaceae	PRU 6211000	Leaves	500
<i>Aspalathus linearis</i>	Rooibos	Leguminosae	PRU 110523	Leaves	125
<i>Barleria albostellata</i>	Grey barleria	Acanthaceae	PRU 096399	Leaves	500
<i>Barleria repens</i>	Small bush violet	Acanthaceae	PRU 081712	Leaves	250
<i>Broussonetia papyrifera</i>	Paper mulberry	Moraceae	PRU 51221	Leaves	500
<i>Buxus macowanii</i>	Cape box	Buxaceae	PRU 110526	Leaves	Na <sup>†</sup>
<i>Carpobrotus edulis</i>	Sour fig	Azioaceae	PRU 096398	Leaves	Na <sup>†</sup>
<i>Ceratonia siliqua</i>	Carob tree	Leguminosae	SM 95502	Leaves	Na <sup>†</sup>
<i>Combretum apiculatum</i>	Red bushwillow	Combretaceae	PRU 110531	Leaves	125
<i>Combretum molle</i>	Velvet bushwillow	Combretaceae	EP 81	Leaves	125
<i>Cotyledon orbiculata</i>	Pig's ear	Crassulaceae	PRU 096402	Leaves	Na <sup>†</sup>
<i>Cryptocarya woodii</i>	Cape laurel	Lauraceae	PRU 064439	Leaves	250
<i>Dahlia imperialis</i>	Dahlia	Asteraceae	PRU 3311010	Leaves	500
<i>Datura stramonium</i>	Thorn apple	Solanaceae	MN 8	Leaves	500
<i>Dichrostachys cinerea</i>	Sickle brush	Leguminosae	PRU 096403	Leaves	500
<i>Diospyros lycioides</i>	Blue brush	Ebenaceae	PRU 118949	Twigs	Na <sup>†</sup>
<i>Dodonaea viscosa</i>	Hopbrush	Sapindaceae	PRU 096404	Leaves	500
<i>Erythrophleum lasianthum</i>	Swazi ordeal tree	Leguminosae	PRU 110525	Leaves	250
<i>Euclea divinorum</i>	Magin gwarra	Ebenaceae	AJ 64	Leaves	250
<i>Euclea natalensis</i>	Natal guarri	Ebenaceae	PRU 95059	Leaves	250
<i>Euclea natalensis</i>	Natal guarri	Ebenaceae	NL 22	Roots	250
<i>Galenia africana</i>	Kraalbos	Aizoaceae	SM 93723	Leaves	125
<i>Gomphocarpus fruticosus</i>	Milkweed	Asclepiadaceae	MN 1	Leaves	250
<i>Greyia flanaganii</i>	Kei bottlebrush	Greyiaceae	P. Van Wyk 2274	Leaves	250
<i>Greyia sutherlandii</i>	Natal bottlebrush	Greyiaceae	PRU 118946	Leaves	125
<i>Harpephyllum caffrum</i>	Wild plum	Anacardiaceae	PRU 118950	Leaves	125
<i>Heteropyxis natalensis</i>	Lavender tree	Myrtaceae	PRU 096405	Leaves	250
<i>Hyaenanche globosa</i>	Hyaena poison	Euphorbiaceae	SM 95499	Leaves	250
<i>Knowltonia vesicatoria</i>	Blister-leaf	Ranunculaceae	PRU 096499	Roots	250
<i>Magnolia grandiflora</i>	Magnolia	Magnoliaceae	PRU 2651000	Leaves	250
<i>Myrsine africana</i>	African boxwood	Myrsinaceae	SM 95503	Stalks	500
<i>Parinari curatellifolia</i>	Mobola plum	Chrysobalanaceae	PRU 096215	Bark	250
<i>Ranunculus repens</i>	Creeping buttercup	Ranunculaceae	PRU 096416	Leaves	125
<i>Rhus lancea</i>	Karee	Anacardiaceae	PRU 110530	Leaves	250
<i>Sclerocarya birrea</i>	Marula	Anacardiaceae	NH 1910	Bark	125
<i>Sideroxylon inerme</i>	White milkwood	Sapotaceae	PRU 96216	Bark	250
<i>Symphytum officinale</i>	Comfrey	Boraginaceae	PRU 096414	Leaves	250
<i>Warburgia salutaris</i>	Pepper-bark tree	Cancellaceae	PRU 110529	Leaves	125
<i>Zanthoxylum capense</i>	Small knob wood	Rutaceae	PRU 096406	Leaves	Na <sup>†</sup>

<sup>†</sup>Na, not active at the highest concentration (500 µg/mL) tested.



**Figure 1:** The DPPH radical scavenging activity of the potential extracts and the positive control (vitamin C): (a) *Aspalathus linearis* ( $EC_{50} = 3.5 \pm 0.5 \mu\text{g/mL}$ ), (b) *Combretum molle* ( $EC_{50} = 9.83 \pm 0.8 \mu\text{g/mL}$ ), (c) *Harpephyllum caffrum* ( $EC_{50} = 2.6 \pm 0.21 \mu\text{g/mL}$ ), (d) *Sclerocarya birrea* ( $EC_{50} = 2.06 \pm 0.03 \mu\text{g/mL}$ ), (e) *Greyia sutherlandii* ( $EC_{50} = 7.9 \pm 0.23 \mu\text{g/mL}$ ), (f) *Combretum apiculatum* ( $EC_{50} = 1.6 \pm 0.02 \mu\text{g/mL}$ ) and (g) vitamin C ( $EC_{50} = 1.98 \pm 0.005 \mu\text{g/mL}$ ).


### Cytotoxicity of selected extracts

Cytotoxicity was assessed on the plant extracts which demonstrated  $EC_{50}$  values of  $\leq 10 \mu\text{g/mL}$  for radical scavenging activity. The plant extracts of *A. linearis*, *G. sutherlandii* and *S. birrea* showed low toxicity with 50% viability of cells ( $EC_{50}$ ) at concentrations of  $125.09 \pm 0.71 \mu\text{g/mL}$ ,  $107.85 \pm 1.53 \mu\text{g/mL}$  and  $92.07 \pm 0.09 \mu\text{g/mL}$ , respectively (Figure 2). During a previous study by our research group, the leaf extract of *H. caffrum* showed toxicity to B16-F10 cells at a concentration of  $100 \mu\text{g/mL}$ .<sup>39</sup> The plant extract of *C. molle* showed moderate toxicity with an  $EC_{50}$  value of  $48.83 \pm 0.21 \mu\text{g/mL}$ , whereas *C. apiculatum* was found to be the most toxic with an  $EC_{50}$  value of  $12.15 \pm 0.03 \mu\text{g/mL}$  and was found to be lethal to almost all cells at the highest concentration of  $400 \mu\text{g/mL}$ . Actinomycin D, the positive control, showed an  $EC_{50}$  value of  $4.5 \times 10^{-3} \pm 0.5 \times 10^{-3} \mu\text{g/mL}$  (Figure 2).

### Discussions

Plant extracts were explored for antibacterial activity against *P. acnes*. Similar to our findings, the ethanolic extract of *Coscinium fenestratum* (Gaertn.) Colebr. (Menispermaceae) inhibited the growth of *P. acnes* at an MIC value of  $46 \mu\text{g/mL}$ .<sup>40</sup> According to Tsai et al.,<sup>41</sup> methanolic extracts of *Rosa damascena* Mill (Rosaceae), *Eucommia ulmoides* Oliv. (Eucommiaceae) and *Ilex paraguariensis* A. St.-Hil. (Aquifoliaceae) inhibited the growth of *P. acnes* at MIC values of  $2000 \mu\text{g/mL}$ ,  $500 \mu\text{g/mL}$  and  $1000 \mu\text{g/mL}$ , respectively. To the best of our knowledge, the present study is the first scientific report of the antibacterial activity of all the selected plants against *P. acnes*. However, some of the plants used in this study have been previously reported to be active against other pathogens. In another study, leaf extracts of *A. linearis* showed zones of inhibition against *Bacillus cereus*, *Micrococcus luteus* and *Candida albicans* of 7.0 mm, 6.4 mm and 8.5 mm, respectively.<sup>42</sup>

The antibacterial activity of *C. apiculatum* against *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli* was reported by Serage.<sup>43</sup> The acetone extract of the stem bark of *C. molle* showed antimicrobial activity against *E. coli* and *Shigella* spp. at an MIC of  $50\ 000 \mu\text{g/mL}$ . The extract also showed inhibitory effects on the fungus *C. albicans* with complete inhibition at a concentration of  $400 \mu\text{g/mL}$ .<sup>44</sup> In a study done by Lining et al.,<sup>45</sup> the crude methanolic extract of *Diospyros lycioides* Desf. (Ebenaceae) showed activity against *Streptococcus mutans* and *Prevotella intermedia* at an MIC of  $1250 \mu\text{g/mL}$ . In contrast, our results showed no activity of the ethanolic extract of *D. lycioides* against *P. acnes*. In another study conducted by Mativandla et al.,<sup>46</sup> the ethanolic extract of *G. africana* showed antimycobacterial activity against *Mycobacterium tuberculosis* and *Mycobacterium smegmatis* at MIC values of  $780 \mu\text{g/mL}$  and  $1200 \mu\text{g/mL}$ , respectively. The ethanolic extract of *H. caffrum* was reported to be active against four bacterial species, namely *Bacillus subtilis*, *E. coli*, *Klebsiella pneumoniae* and *S. aureus*, while an aqueous extract showed activity against *C. albicans*.<sup>47</sup> The acetone extracts of the bark and leaves of *S. birrea* were reported to be active against *S. aureus*, *P. aeruginosa*, *E. coli* and *Enterococcus faecalis* at MIC values ranging from  $150 \mu\text{g/mL}$  to  $3000 \mu\text{g/mL}$ .<sup>48</sup> In a study done by Motsei et al.,<sup>49</sup> the leaf extracts of *W. salutaris* inhibited growth of *C. albicans* at MIC values ranging from  $12\ 500 \mu\text{g/mL}$  to  $25\ 000 \mu\text{g/mL}$  and the bark extracts showed growth of inhibition against *S. aureus*, *Staphylococcus epidermis*, *B. subtilis* and *E. coli*.<sup>50</sup> No reports regarding the antimicrobial activity of *G. sutherlandii* and *R. repens* were found in the literature. However, in the present study, both of these plants showed growth inhibitory activity against *P. acnes* at an MIC of  $125 \mu\text{g/mL}$ . In a study conducted by Eloff and Katerere,<sup>51</sup> the acetone and chloroform leaf extracts of *A. galpinii* inhibited the growth of *S. aureus* and *E. coli*. Similar to our findings, the ethanol bark extract of



**Figure 2:** The cytotoxic effects of the plant extracts and the positive control (actinomycin D) on mouse melanocytes B16-F10: (a) *Aspalathus linearis* ( $EC_{50} = 125.09 \pm 0.71 \mu\text{g/mL}$ ), (b) *Greyia sutherlandii* ( $EC_{50} = 107.85 \pm 1.53 \mu\text{g/mL}$ ), (c) *Sclerocarya birrea* ( $EC_{50} = 92.07 \pm 0.09 \mu\text{g/mL}$ ) and (d) actinomycin D ( $EC_{50} = 4.5 \times 10^{-3} \pm 0.5 \times 10^{-3} \mu\text{g/mL}$ ).

*A. galpinii* exhibited good inhibitory effect on *P. acnes* (MIC 62.5 µg/mL). However, no reports on the antimicrobial activity of bark extracts of *A. galpinii* were found in the literature.

Acne is associated with the production of free radicals along with the infection of *P. acnes*. Reactive oxygen species are produced as a result of the action of hydrolytic enzymes released from bacteria on the follicular walls of pilosebaceous units. Therefore, the plant extracts were evaluated for antioxidant activity along with antibacterial activity. In our study, the ethanol extracts of *A. linearis*, *C. apiculatum*, *H. caffrum*, *S. birrea*, *C. molle* and *G. sutherlandii* exhibited significant antioxidant activity with EC<sub>50</sub> values of ≤10 µg/mL. Our results are in agreement with other researchers. During a previous study by Joubert et al.<sup>52</sup>, the DPPH radical scavenging activity of *A. linearis* and its constituents were confirmed. The polar fractions of *C. apiculatum* showed antioxidant activity with an EC<sub>50</sub> value of 3.91 µg/mL.<sup>53</sup> The DPPH radical scavenging activity of *H. caffrum* and *S. birrea* was confirmed by Moyo et al.<sup>54</sup> with EC<sub>50</sub> values of 6.8 µg/mL and 5.02 µg/mL, respectively. In another study, acetone and dichloromethane extracts of *C. molle* displayed antioxidant activity after spraying with DPPH.<sup>55</sup> It has been reported that DPPH free radicals abstract the phenolic hydrogen of the electron-donating molecule, which could be the general mechanism for the scavenging action of flavonoids.<sup>56</sup> Based on the mechanism of reduction of the DPPH molecule that is correlated with the presence of hydroxyl groups on the antioxidant molecule, the antioxidant activity of the polar plant extracts in the present study can be explained as a result of the presence of their phytoconstituents (phenolics or flavonoids) which are radical scavengers with an available hydroxyl group and are known to occur abundantly in plant species.

In order to evaluate the therapeutic potential of the plants, the cytotoxicity of selected samples was tested on B16-F10 cells. To the best of our knowledge, the cytotoxicity of the extracts described in the present study is reported for the first time. However, previous researchers have documented similar cytotoxic effects on different cell lines. In a study by McGaw et al.<sup>57</sup>, *A. linearis* showed low toxicity on vero cells and brine shrimp larvae with LD<sub>50</sub> values of >1000 µg/mL. *S. birrea* showed low cytotoxicity on vero cells with an IC<sub>50</sub> value of 361.24 µg/mL.<sup>58</sup> According to previous studies by Fyhrquist et al.<sup>59</sup> on the cytotoxicity of *C. molle*, the extract showed IC<sub>50</sub> values of 27.7 µg/mL, 72.6 µg/mL and 42.6 µg/mL on T24 (bladder carcinoma), HeLa (cervical carcinoma) and MCF-7 (breast carcinoma) cells, respectively, while the *C. apiculatum* extract showed IC<sub>50</sub> values of 65.0 µg/mL and 40.1 µg/mL for T24 and MCF-7 cells, respectively. No records of cell cytotoxicity for *G. sutherlandii* were found in the literature.

The results shown in this study prove the capability of medicinal plants as anti-acne agents, although the mode of action and in vivo studies are required to give conclusive results.

## Conclusions

Based on the results obtained, it can be concluded that the ethanol bark extract of *A. galpinii* demonstrated the best activity against *P. acnes* with acceptable antioxidant activity. This plant might have other attributes that were not investigated in the present study which could be useful in the treatment of *P. acnes*. Although the plant extracts of *H. caffrum*, *C. apiculatum* and *C. molle* showed good antibacterial and excellent antioxidant activity, these samples also showed moderate toxicity to mouse melanocyte cells. The plant extracts of *A. linearis*, *S. birrea* and *G. sutherlandii* also exhibited good antibacterial and antioxidant activity but had low toxicity to the mouse melanocytes; these extracts therefore have potential as anti-acne agents, either alone or in combination.

## Acknowledgements

We thank the University of Pretoria and the National Research Foundation (South Africa) for financial grants.

## Authors' contributions

R.S. conducted the experiments and drafted the manuscript; N.L. supervised the work and edited the manuscript.

## References

1. Bloch B. Metabolism, endocrine glands and skin disease, with special reference to acne vulgaris and xanthoma. *Brit J Dermatol.* 1931;43:61–87. <http://dx.doi.org/10.1111/j.1365-2133.1931.tb09468.x>
2. Munro-Ashman D. Acne vulgaris in public schools. *Trans St John's Hosp Dermatol Soc.* 1963;49:144–148.
3. Burton JL, Cunliffe WJ, Stafford I, Shuster S. The prevalence of acne vulgaris in adolescence. *Brit J Dermatol.* 1971;85:119–126. <http://dx.doi.org/10.1111/j.1365-2133.1971.tb07195.x>
4. Rademaker M, Garioch JJ, Simpson NB. Acne in school children: No longer a concern for dermatologists. *Brit Med J.* 1989;298:1217–1219. <http://dx.doi.org/10.1136/bmj.298.6682.1217>
5. Shaw L, Kennedy C. The treatment of acne. *Paediatr Child Health.* 2007;17(10):385–389. <http://dx.doi.org/10.1016/j.paed.2007.07.005>
6. Arican O, Kurutas EB, Sasmaz S. Oxidative stress in patients with acne vulgaris. *Mediat Inflamm.* 2005;6:380–384. <http://dx.doi.org/10.1155/MI.2005.380>
7. Hutchings A, Scott AH, Lewis G, Cunningham AB. Zulu medicinal plants. An inventory. Pietermaritzburg: University of Natal Press; 1996.
8. Van Wyk BE, Gerick N. People's plants. A guide to useful plants of southern Africa. Pretoria: Briza Publications; 2006.
9. Kokwaro O. Medicinal plant of East Africa. Kampala/Nairobi/Dar es Salaam: East African Literature Bureau; 1976.
10. Davis RH, Rosental KY, Cesario LR, Rouw GA. Processed *Aloe vera* administered topically inhibits inflammation. *J Am Podiat Med.* 1989;79:395–397. <http://dx.doi.org/10.7547/87507315-79-8-395>
11. Lee KY, Weintraub ST, Yu BP. Isolation and identification of a phenolic antioxidant from *Aloe barbadensis*. *Free Radical Bio Med.* 2000;28(2):261–265. [http://dx.doi.org/10.1016/S0891-5849\(99\)00235-X](http://dx.doi.org/10.1016/S0891-5849(99)00235-X)
12. Van Wyk BE, Van Oudtshoorn B, Gericke N. Medicinal Plants of South Africa. Pretoria: Briza Publications; 2007.
13. Wink M, Van Wyk BE. Mind-altering and poisonous plants of the world. Pretoria: Briza Publications; 2008.
14. Agroforestry Tree Database [database on the Internet]. No date [cited 2010 Nov 27]. Available from: [http://www.worldagroforestry.org/treedb/AFTPDFS/Annona\\_senegalensis.pdf](http://www.worldagroforestry.org/treedb/AFTPDFS/Annona_senegalensis.pdf)
15. Ziyat A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M, Benjelloun W. Phytotherapy of hypertension and diabetes in oriental Morocco. *J Ethnopharmacol.* 1997;58:45–54. [http://dx.doi.org/10.1016/S0378-8741\(97\)00077-9](http://dx.doi.org/10.1016/S0378-8741(97)00077-9)
16. Ziyat A, Boussairi EH. Cardiovascular effects of *Arbutus unedo* L. in spontaneously hypertensive rats. *Phytother Res.* 1998;12:110–113. [http://dx.doi.org/10.1002/\(SICI\)1099-1573\(199803\)12:2<110::AID-PTR199>3.0.CO;2-5](http://dx.doi.org/10.1002/(SICI)1099-1573(199803)12:2<110::AID-PTR199>3.0.CO;2-5)
17. Kivcak B, Mert T. Quantitative determination of tocopherol in *Arbutus unedo* by TLC-densitometry and colorimetry. *Fitoterapia.* 2001;72:656–661. [http://dx.doi.org/10.1016/S0367-326X\(01\)00305-7](http://dx.doi.org/10.1016/S0367-326X(01)00305-7)
18. Pabuccuoglu A, Kivcak B, Bas M, Mert T. Antioxidant activity of *Arbutus unedo* leaves. *Fitoterapia.* 2003;74:597–599. [http://dx.doi.org/10.1016/S0367-326X\(03\)00110-2](http://dx.doi.org/10.1016/S0367-326X(03)00110-2)
19. Stephen AO. Micro propagation and medicinal properties of *Barleria greenii* and *Huernia hystrix* [PhD thesis]. Pietermaritzburg: University of KwaZulu-Natal; 2010.
20. Whistler WA, Elevitch CR. *Broussonetia papyrifera* (paper mulberry). Species profiles for Pacific Island Agroforestry [document on the Internet]. c2006 [cited 2010 Dec 02]. Available from: <http://www.agroforestry.net/tii/Broussonetia-papermulb.pdf>
21. Kivcak B, Mert T, Ozturk HT. Antimicrobial and cytotoxic activities of *Ceratonia siliqua* L. extracts. *Turk J Biol.* 2002;26:197–200.
22. Mbambezeli G. *Cryptocarya wodii* Engl. [homepage on the Internet]. c2005 [cited 2010 Nov 27]. Available from: <http://www.plantzfrica.com/plantcd/cryptocarwood.htm>

23. Roberts M. *Dahlia*. Edible and medicinal flowers. Cape Town: Spearhead Press; 2007.
24. Vries FA, Bitar HEL, Green IR, Klassen JA, Mabusela WT, Bodo B, et al. An antifungal extract from the aerial parts of *Galenia africana*. In: Proceedings of the 11th NAPRECA Symposium; 2005 August 9–12; Antananarivo, Madagascar. Nairobi: NAPRECA; 2005. p. 123–131. Available from: <http://napreca.net/publications/11symposium/>
25. Mbambezeli G. *Greyia flanaganii* Bolus [homepage on the Internet]. c2002 [cited 2010 Nov 27]. Available from: <http://www.plantzafrica.com/plantefg/greyiaflan.htm>.
26. Martinez M. Las plantas medicinales de Mexico [Medicinal plants of Mexico]. 4th ed. Mexico City: Ediciones Botas; 1959. p. 343–347. Spanish.
27. Mellado V, Chavez SMA, Lozoya X. Pharmacological screening of the aqueous extracts of *Magnolia grandiflora* L. Arch Med Res. 1980;11(3):335–346.
28. Beentje HJ. Kenya trees, shrubs and lianas. Nairobi: National Museums of Kenya; 1994.
29. Mabogo DEN. The ethnobotany of Vhavenda [MSc thesis]. Pretoria: University of Pretoria; 1990.
30. Maharaj V, Glen HF. *Parinari curatifolia* [homepage on the Internet]. c2008 [cited 2010 Nov 27]. Available from: <http://www.plantzafrica.com/plantnop/parinari curat.htm>.
31. Moreman D. Native American ethnobotany. Portland, OR: Timber Press; 1998.
32. Gundidza M, Gweru N, Mmbengwa V, Ramalivhana NJ, Magwa Z, Samie A. Phytoconstituents and biological activities of essential oil from *Rhus lancea* L.F. Afr J Biotechnol. 2008;7(16):2787–2789.
33. Bosman F. *Sideroxylon inerme* L. [homepage on the Internet]. c2006 [cited 2010 Nov 27]. Available from: <http://www.plantzafrica.com/plantqrs/sideroxinerm.htm>
34. Buchman DD. Herbal medicine. New York: Gramercy Books/Random House Value Publishing; 1988. p. 101–171.
35. Smith A. A contribution to South African Materia Medica, chiefly from plants in use among the natives. Cape Town: Juta; 1985.
36. Mapunya MB, Hussein AA, Rodriguez B, Lall N. Tyrosinase activity of *Greyia flanaganii* Bolus constituents. Phytomedicine. 2011;18:1006–1012. <http://dx.doi.org/10.1016/j.phymed.2011.03.013>
37. Du Toit R, Volsteed Y, Apostolides Z. Comparison of the antioxidant content of the fruits, vegetables and teas measured as vitamin C equivalents. Toxicology. 2001;166:63–69. [http://dx.doi.org/10.1016/S0300-483X\(01\)00446-2](http://dx.doi.org/10.1016/S0300-483X(01)00446-2)
38. Kuete V. Potential of Cameroonian plants and derived-products against microbial infections. Planta Med. 2010;76:1479–1491. <http://dx.doi.org/10.1055/s-0030-1250027>
39. Mapunya MB, Nikolova RV, Lall N. Melanogenesis and antityrosinase activity of selected South African plants. Evid Based Complement Alternat Med. 2012, Article ID 374017, 6 pages. <http://dx.doi.org/10.1155/2012/374017>
40. Kumar GS, Jayaveera KN, Kumar CK, Sanjay UP, Swamy BM, Kumar DV. Antimicrobial effects of Indian medicinal plants against acne-inducing bacteria. Trop J Pharm Res. 2007;6(2):717–723. <http://dx.doi.org/10.4314/tjpr.v6i2.14651>
41. Tsai T, Tsai T, Wu W, Tseng JT, Tsai P. In vitro antimicrobial and anti-inflammatory effects of herbs against *Propionibacterium acnes*. Food Chem. 2010;119(3):964–968. <http://dx.doi.org/10.1016/j.foodchem.2009.07.062>
42. Almajano MP, Carbo R, Jimenez JAL, Gordon MH. Antioxidant and antimicrobial activities of tea infusions. Food Chem. 2008;108(1):55–63. <http://dx.doi.org/10.1016/j.foodchem.2007.10.040>
43. Serage A. Isolation and characterization of antibacterial compounds present in *Combretum apiculatum* subsp. *apiculatum* [MSc thesis]. Pretoria: Programme for Phytomedicine, University of Pretoria; 2008.
44. Asres K, Mazumder A, Bucar F. Antibacterial and antifungal activities of *Combretum molle*. Ethop Med J. 2006;44:269–277.
45. Lining C, Guo-Xian W, Van der Bijl P, Wu CD. Namibian chewing stick, *Diospyros lycioides*, contains antibacterial compounds against oral pathogens. J Agri Food Chem. 2000;48(3):909–914. <http://dx.doi.org/10.1021/jf9909914>
46. Mativandele SPN, Meyer JMM, Hussein AA, Houghton PJ, Hamilton CJ, Lall N. Activity against *Mycobacterium smegmatis* and *M. tuberculosis* by extracts of South African medicinal plants. Phytother Res. 2008;22(6):841–845. <http://dx.doi.org/10.1002/ptr.2378>
47. Buwa LV, Van Staden J. Effects of collection time on the antimicrobial activities of *Harpephyllum caffrum* bark. S Afr J Bot. 2007;73(2):242–247. <http://dx.doi.org/10.1016/j.sajb.2006.09.006>
48. Eloff JN. Antibacterial activity of Marula (*Sclerocarya birrea* (A. Rich.) Hochst. subsp. *caffra* (Sond.) Kokwaro, (Anacardiaceae) bark and leaves. J Ethnopharmacol. 2001;76(3):305–308. [http://dx.doi.org/10.1016/S0378-8741\(01\)00260-4](http://dx.doi.org/10.1016/S0378-8741(01)00260-4)
49. Motsei ML, Lindsey KL, Van Staden J, Jager AK. Screening of traditionally used South African plants for antifungal activity against *Candida albicans*. J Ethnopharmacol. 2003;86(2):235–241. [http://dx.doi.org/10.1016/S0378-8741\(03\)00082-5](http://dx.doi.org/10.1016/S0378-8741(03)00082-5)
50. Rabe T, Van Staden J. Antibacterial activity of South African plants used for medicinal purposes. J Ethnopharmacol. 1997;56(1):81–87. [http://dx.doi.org/10.1016/S0378-8741\(96\)01515-2](http://dx.doi.org/10.1016/S0378-8741(96)01515-2)
51. Eloff JN, Katerere DR. Variation in chemical composition, antibacterial and antioxidant activity of fresh and dried *Acacia* leaf extracts. S Afr J Bot. 2004;70(2):303–305.
52. Joubert E, Winterton P, Britz TJ, Ferreira D. Superoxide anion and  $\alpha,\alpha$ -diphenyl- $\beta$ -picrylhydrazyl radical scavenging capacity of rooibos (*Aspalathus linearis*) aqueous extracts, crude phenolic fractions, tannin and flavonoids. Food Res Int. 2004;37(2):133–138. <http://dx.doi.org/10.1016/j.foodres.2003.09.011>
53. Aderogba MA, Kgate DT, McGaw LJ, Eloff JN. Isolation of antioxidant constituents from *Combretum apiculatum* subsp. *apiculatum*. S Afr J Bot. 2012;79:125–131. <http://dx.doi.org/10.1016/j.sajb.2011.10.004>
54. Moyo M, Ndhkala AR, Finnie JF, Van Staden J. Phenolic composition, antioxidant and acetylcholinesterase inhibitory activities of *Sclerocarya birrea* and *Harpephyllum caffrum* (Anacardiaceae) extracts. Food Chem. 2010;123(1):69–76. <http://dx.doi.org/10.1016/j.foodchem.2010.03.130>
55. Masoko P, Eloff JN. Screening of twenty-four South African *Combretum* and six *Terminalia* species (Combretaceae) for antioxidant activities. Afr J Tradit Complement Altern Med. 2007;4(2):231–239.
56. Ratty AK, Sunamoto J, Das NP. Interaction of flavonols with 1,1 – diphenyl- 2-picrylhydrazyl free radical liposomal membranes and soybean lipoxigenase-1. Biochem Pharmacol. 1998;37:989–995. [http://dx.doi.org/10.1016/0006-2952\(88\)90499-6](http://dx.doi.org/10.1016/0006-2952(88)90499-6)
57. McGaw LJ, Steenkamp V, Eloff JN. Evaluation of Athrixia bush tea for cytotoxicity, antioxidant activity, caffeine content and presence of pyrrolizidine alkaloids. J Ethnopharmacol. 2007;110(1):16–22. <http://dx.doi.org/10.1016/j.jep.2006.08.029>
58. Gathirwa JW, Rukunga GM, Njagi ENM, Omar SA, Mwitari PG, Guantai AN, et al. The in vitro anti-plasmodial and in vivo anti-malarial efficacy of combinations of some medicinal plants used traditionally for treatment of malaria by the Meru community in Kenya. J Ethnopharmacol. 2008;115(2):223–231. <http://dx.doi.org/10.1016/j.jep.2007.09.021>
59. Fyhrquist P, Mwasumbi L, Vuorela P, Vuorela H, Hiltunen R, Murphy C, et al. Preliminary antiproliferative effects of some species of *Terminalia*, *Combretum*, and *Pteleopsis* collected in Tanzania on some human cancer cell lines. Fitoterapia. 2006;77(5):358–366. <http://dx.doi.org/10.1016/j.fitote.2006.05.017>

