

Hereditary haemorrhagic telangiectasia in North African and sub-Saharan patients

To the Editor: Hereditary haemorrhagic telangiectasia (HHT) or Osler-Weber-Rendu disease is an autosomal-dominant inherited vascular disease, characterised by the presence of mucocutaneous telangiectasia and visceral arteriovenous malformations (AVMs). Three main causative genes are known: *ENG*, *ACVRL1* and *SMAD4*. *BMP9* has also been shown to be involved in a small number of cases. We report two cases of HHT in North African and sub-Saharan patients.

HHT is a vascular dysplasia that affects approximately 1 in 5 000 - 8 000 individuals,^[1] with regional differences due to founder effects.^[2,3] The clinical diagnosis of HHT is based on the Curaçao criteria,^[1] including: spontaneous and recurrent nosebleeds; mucocutaneous telangiectasia at characteristic sites (nose, lips, oral cavity, fingertips and gastrointestinal mucosa); AVMs in lungs, liver, gastrointestinal tract, brain and spinal cord; and a first-degree relative with HHT.

Approximately 85% of HHT patients^[3] have a mutation identified in either *ENG* (OMIM#187300, HHT1)^[4] or *ACVRL1* (OMIM#600376, HHT2),^[5] and 2% of the HHT families have a mutation identified in the *SMAD4* gene, present in HHT in association with juvenile polyposis (OMIM #175050, JPHT).^[6]

For both *ENG* and *ACVRL1*, many mutations have been reported by independent groups in different countries, in most cases without any evidence of a founder effect, which could be suspected only in a limited number of cases.^[2,7] As is known, HHT in people of black African descent is a rarity. To date, only one paper reports a mutation (exon 7 of *ACVRL1*: c.818T>C, p.L273P) in an HHT black adult patient born in Kenya,^[8] as Westermann *et al.*^[9] discuss in their letter. The other cases in literature were published before the identification of the two disease-related genes.^[2] A December 2012 census showed that the number of immigrants from North Africa and sub-Saharan countries, residing regularly in Italy, for whom the public healthcare service is fully available, is 760 500 and 350 000, respectively. However, a high number of non-registered immigrants are certainly




Fig. 1. SpliceAid analysis of *ENG* mutation (c. 220-1G>C, splice defect). (A) *hnRNP* exonic splicing silencer (ESS) in wild-type sequence of intron2/exon3 of *ENG* (circled); (B) *SRp40* exonic splicing enhancer (ESE) in mutated sequence of intron2/exon3 of *ENG* (circled).

present. The countries of origin of the two patients are those with the highest number of immigrants to Italy from North African or sub-Saharan areas – Morocco and Senegal.

We wish to report our observation on two patients among the ~500 index cases, collected by the centres caring for HHT in northern Italy, in Pavia and Crema. The HHT diagnosis was arrived at according to Curaçao criteria. The first case is a patient born in Morocco (regular immigrants: ~501 600) with a 'possible' diagnosis of HHT (2/4 Curaçao criteria). The second patient was born in Senegal (regular immigrants: ~81 000) with a 'definite' diagnosis of HHT (3/4 Curaçao criteria).

Molecular analysis of *ENG* and *ACVRL1*, according to Olivieri *et al.*,^[10] disclosed a mutation in intron 2 of *ENG* (*c.220-1G>C*, splice defect) and a mutation in exon 8 of *ACVRL1* (*c.1232G>A*, p.R411Q), respectively. Analysis of exon 3 of *ENG*, including intron-exon boundaries, covers 20% of the mutations identified in this gene in our lab. The mutation observed is yet unpublished. Analysis of splice sites with SpliceAid (<http://www.introni.it/>) (Fig. 1) and Human Splicing Finder (<http://www.umd.be/HSF/>), bioinformatics tools useful for predicting the effect of splicing mutations, shows the replacement of an exonic splicing silencer (ESS) with an exonic splicing enhancer (ESE). Overall, bioinformatics tests suggest that the mutation is pathogenetic. Exon 8 of the *ACVRL1* gene carries 24% of the mutations identified in this gene in our lab. We observed the same mutation in 5/38 index cases from unrelated families and it has also been reported several times in the HHT mutation database (<http://arup.utah.edu/database/hht/>) in patients from different ethnic origins.

Overall, these data support the observation by Westermann *et al.*^[9] regarding the rarity of HHT in patients from Africa. We are aware of the limitations associated with reporting single cases, but we feel a short report of this observation is relevant as data are totally lacking regarding the incidence and prevalence of HHT, and the relative involvement of *ENG* or *ACVRL1* in the African population.

Cecilia Canzonieri

Federica Ornati

General Biology and Medical Genetics, Department of Molecular Medicine,
University of Pavia, Italy
cecilia.canzonieri@gmail.com

Elina Matti

Francesco Chu

Department of Otorhinolaryngology, Foundation IRCCS, San Matteo Polyclinic,
Pavia, Italy

Guido Manfredi

Department of Gastroenterology, Maggiore Hospital, Crema, Italy

Carla Olivieri

General Biology and Medical Genetics, Department of Molecular Medicine,
University of Pavia, Italy

Elisabetta Buscarini

Department of Gastroenterology, Maggiore Hospital, Crema, Italy

Fabio Pagella

Department of Otorhinolaryngology, Foundation IRCCS, San Matteo Polyclinic,
Pavia, Italy

- Westermann CJ, Rosina AF, de Vries V, de Coteau PA. The prevalence and manifestations of hereditary hemorrhagic telangiectasia in the Afro-Caribbean population of the Netherlands Antilles: A family screening. *Am J Med Genet A* 2003;116A(4):324-328. [<http://dx.doi.org/10.1002/ajmg.a.10002>]
- Lesca G, Plauchu H, Coulet F, et al. Molecular screening of *ALK1/ACVRL1* and *ENG* genes in hereditary hemorrhagic telangiectasia in France. *Hum Mutat* 2004;23(4):289-299. [<http://dx.doi.org/10.1002/humu.20017>]
- McAllister KA, Baldwin MA, Thukkani AK, et al. Six novel mutations in the endoglin gene in hereditary hemorrhagic telangiectasia type 1 suggest a dominant-negative effect of receptor function. *Hum Mol Genet* 1995;4(10):1983-1985.
- Johnson DW, Berg JN, Baldwin MA, et al. Mutations in the activin receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia type 2. *Nat Genet* 1996;13(2):189-195. [<http://dx.doi.org/10.1038/ng0696-189>]
- Gallione CJ, Repetto GM, Legius E, et al. A combined syndrome of juvenile polyposis and hereditary haemorrhagic telangiectasia associated with mutations in *MADH4* (*SMAD4*). *Lancet* 2004;363(9412):852-859. [[http://dx.doi.org/10.1016/S0140-6736\(04\)15732-2](http://dx.doi.org/10.1016/S0140-6736(04)15732-2)]
- Lesca G, Olivieri C, Burnichon N, et al. Genotype-phenotype correlations in hereditary hemorrhagic telangiectasia: Data from the French-Italian HHT network. *Genet Med* 2007;9(1):14-22. [<http://dx.doi.org/10.1097/GIM.0b013e31802d8373>]
- Kitonyi GW, Wambugu PM, Oburra HO, Ileri JM. Hereditary haemorrhagic telangiectasia in a black adult male: Case report. *East Afr Med J* 2008;85(8):412-416.
- Westermann CJ, Kitonyi GW, Letteboer TG. Is hereditary haemorrhagic telangiectasia rare in the black race? The first sub-Saharan mutation. *Haemophilia* 2011;17(1):e244. [<http://dx.doi.org/10.1111/j.1365-2516.2010.02353.x>]
- Olivieri C, Pagella F, Semino L, et al. Analysis of *ENG* and *ACVRL1* genes in 137 HHT Italian families identifies 76 different mutations (24 novel). Comparison with other European studies. *J Hum Genet* 2007;52(10):820-829. [<http://dx.doi.org/10.1007/s10038-007-0187-5>]

S Afr Med J 2014;104(4):256-257. DOI:10.7196/SAMJ.7816

1. Faughnan ME, Palda VA, Garcia-Tsao G, et al. International guidelines for the diagnosis and management of hereditary haemorrhagic telangiectasia. *J Med Genet* 2011;48(2):73-87. [<http://dx.doi.org/10.1136/jmg.2009.069013>]