



CLINICAL IMAGES

Cutaneous bacillary angiomatosis: A marker of systemic disease in HIV

N Grilo, D Modi, P Barrow

A 33-year-old man presented with a 5-week history of relapsing fever, weight loss and right upper quadrant pain. He was pyrexial, the liver and spleen were enlarged and tender, and he had a few pyogenic granuloma-like tumours on the skin (Fig. 1). He was HIV positive (CD4 cell count 119/ μ l), and had raised liver enzyme levels (alkaline phosphatase (ALP) 258 IU/l and gamma-glutamyl transpeptidase (GGT) 218 IU/l). Results of hepatitis studies and the alpha-fetoprotein level were normal, blood cultures were negative and a chest radiograph was normal. An abdominal ultrasound scan showed an enlarged and markedly echogenic liver and spleen with multiple hypo-echoic foci in keeping with possible micro-abscesses. The skin and liver showed similar histological features: clusters of purple, granular material that stained positive for organisms, confirmed on the Warthin-Starry stain (Figs 2a, 2b and 2c).

The polymerase chain reaction (PCR) confirmed the presence of *Bartonella* spp. on both biopsy specimens. The skin lesions had almost completely disappeared after 1 month on erythromycin 500 mg 4 times a day, but the patient was subsequently lost to follow-up.

Discussion

Bacillary angiomatosis (BA) is caused by the Gram-negative bacteria *B. henselae* and *B. quintana*. Cutaneous BA was first described in 1983, and the first case in South Africa (where disease prevalence in the host, the domestic cat, is 24%)¹ was reported in 1993.² BA is difficult to diagnose, requiring culture for at least 21 days; serological studies are often unreliable, and special staining with the Warthin-Starry stain is used to confirm the tissue diagnosis. The prevalence of *Bartonella*



Fig. 1. Pyogenic granuloma-like lesion above right ear




Fig. 2a. Low-power H&E stain of the lesion.

Dr Nella Grilo qualified and did her postgraduate training at the University of the Witwatersrand. She is currently a consultant dermatologist at Chris Hani Baragwanath Hospital, Johannesburg. Professor Deepak Modi also trained at Wits, and is currently academic head of the Division of Dermatology there. Dr Peter Barrow is a gastroenterologist, currently working in Dubai.

Corresponding author: N Grilo (27823189392@vodamail.co.za)




Fig. 2b. Higher-power H&E stain showing purple, granular material.




Fig. 2c. Positive Warthin-Starry stain demonstrating the organisms.

bacteraemia (nested PCR) was 10% at the Johannesburg HIV outpatient clinics.³ The treatment of choice is erythromycin 500 mg 4 times a day for 3 months; also useful are doxycycline, ceftriaxone and the fluoroquinolones.

Examination of the skin is mandatory because cutaneous lesions may be a marker of systemic BA infection, especially in HIV-positive patients, and skin biopsy is much safer than liver biopsy for tissue diagnosis.

1. Kelly PJ, Matthewman LA, Hayter D, et al. *Bartonella (Rochalimaea) henselae* in southern Africa – evidence for infections in domestic cats and implications for veterinarians. *J S Afr Vet Assoc* 1996; 67: 182-187.
2. Levy GR, Nayler S. Bacillary angiomatosis. The first case reported in South Africa. *S Afr Med J* 1993; 83: 855-856.
3. Frean J, Arndt S, Spence D. High rate of *Bartonella henselae* infection in HIV-positive outpatients in Johannesburg, South Africa. *Trans R Soc Trop Med Hyg* 2002; 96: 549-550.



Need Healthcare -
that does not cost
an arm and a leg?



RESOLUTION
health

Tel: 0861 11 RESO / 7376
www.resomed.co.za