

---

# CLINICAL ARTICLE

---

## Efficacy of an exercise programme on the functional capacity and disease activity in females with rheumatoid arthritis

Dr DC Janse van Rensburg\* MBChB, MMed, MSc

Dr L Fletcher<sup>‡</sup> PhD

Prof M Viljoen\* PhD, PhD

Ms C Coertzen<sup>‡</sup> BA(MBK) Hons

Mrs CC Grant\* MSc

Dr DA Ramagole\* MBChB, MSc

Dr RM Collins\* MBChB, MSc

Prof JA Ker\* MBChB, MMed, MD

University of Pretoria

\*Section Sports Medicine <sup>‡</sup>Department of Statistics <sup>‡</sup>Department of Physiology  
<sup>‡</sup>Department of Biokinetics, Sport and Leisure Sciences <sup>‡</sup>Faculty of Health Sciences

### Correspondence:

Dr DC Janse van Rensburg

Section Sports Medicine

University of Pretoria

P O Box 12651

0028 Hatfield

Tel: (012) 420-6057

Fax: (012) 362-3369

E-mail: vrensburg@sport.up.ac.za

### Abstract

#### Background

Rheumatoid arthritis (RA) is a chronic, immune-inflammatory disease of unknown aetiology affecting the synovial membrane of joints and surrounding tissues. Typically RA affects both large and small joints in a bilateral, symmetrical, poly-articular fashion. Degradation of bone, cartilage and muscle eventually lead to a reduction in physical function. The purpose of this study was to determine the efficacy of an endurance exercise programme on the fitness parameters (flexibility, strength and aerobic measurements), quality of life (visual analogue scale and health assessment questionnaire) and disease activity (DAS28[4] version with CRP) of female RA patients.

### Methods

Female RA patients were randomly allocated to the experimental group (EG) (n=19) and the control group (CG) (n=8). All participants went through a battery of tests before the intervention, and again after completion of the study. The 12 week training programme consisted of three 45-minute training sessions per week and included walking or aquatics, as well as stretches and isotonic strengthening exercises. The Mann-Whitney U test was used to compare measurements between groups. The Wilcoxon signed-rank test was used to compare baseline and post-intervention measurements within each group.

### Results

At the initiation of the study the CG and the EG were comparable for fitness, quality of life and disease activity. On completion of the training programme, statistically significant improvements at the 5% level of significance were seen between the EG and CG, in favour of the EG, for left lateral flexion (p=0.015) and the 1 mile walk test (p=0.011). Within the EG there were improvement of knee flexion left (p=0.026), knee extension (right p=0.011; left p=0.009), scratch test (right p=0.007; left p=0.01), chair sit and reach (right p=0.011; left p<0.001), strength parameters (p<0.05), 1 mile walk test (p<0.001), VO<sub>2</sub> max (p=0.01) and DAS scores (p<0.001). Within the CG, improvements were shown for knee extension (right p=0.05; left p=0.013). Although their strength parameters improved significantly it was not in the same order as for the EG. The CG had a decline in their aerobic measurements but their HAQ score improved (p=0.03).

### Conclusion

An endurance exercise programme, combined with isotonic strengthening exercise and stretching, improves the functional capacity, quality of life and disease activity of female patients with RA. Attention received during the study may have led to some placebo-induced improvements in control subjects, but not to the same extent as those involved in exercise programmes.

## Introduction

Rheumatoid arthritis (RA) is a chronic, immune-inflammatory disease of unknown aetiology affecting the synovial membrane of joints and surrounding tissues.<sup>1-4</sup> Typically RA affects both large and small joints in a bilaterally symmetrical polyarticular fashion. The most common joints involved are the small joints of the hands and feet followed by the wrists and ankles. The affected joints are swollen, warm, tender and painful on movement.<sup>1</sup>

RA causes various physical impairments in those affected by the disease.<sup>5</sup> Some of these may include: an inhibition of muscle contraction as result of joint effusion; myositis; muscle atrophy due to decreased activity levels which leads to a decrease in muscle strength; a loss of joint motion; and reduced aerobic capacity secondary to systemic effects and reduced activity levels. Persons with RA, especially those severely affected by the disease, are in general 33% to 55% weaker than their healthy counterparts.<sup>6</sup>

Although there is no cure for RA, much can be done to manage the condition. Four major treatment approaches are recognised in the management of RA, including medication, physical exercise, joint protection and lifestyle changes, and surgical intervention.<sup>7</sup>

According to the American College of Sports Medicine the primary objectives of exercise therapy in patients with RA is (1) to preserve or restore range of motion (ROM) and flexibility around affected joint(s); (2) to increase muscle strength and endurance to build joint stability; and (3) to increase aerobic capacity in order to enhance psychological state and decrease the risk of cardiovascular disease.<sup>2</sup>

A comprehensive exercise programme for RA patients is said to include aerobic exercise at a moderate intensity for three to five days a week, isometric or isotonic strength training exercises three days a week, as well as stretching exercises at least once daily.<sup>8</sup> Several papers suggest that exercise may improve functional capacity (i.e. quality of life and fitness parameters),<sup>9-12</sup> and have an effect on disease activity.<sup>13-17</sup>

The objective of this study was to measure the effect of an endurance training programme on the fitness parameters, quality of life and disease activity of females suffering from RA.

## Materials and methods

### Study design

The study was conducted at the University of Pretoria, and ethical clearance was obtained from the Ethical Committee, Faculty of Health Sciences. All participants signed an informed consent document. A prospective analytical pre-post group comparison was adopted.

**Persons with RA are in general 33% to 55% weaker than their healthy counterparts**

Participants were selected based on the inclusion and exclusion criteria (Table I).

A random sample, equally allocated to a control group (CG) and an experimental group (EG) consisting of two subsets (Aquatic exercise and Land exercise) was planned. The final sample consisted of 37 patients who were randomly allocated to the EG (n=25) and the CG (n=12) by drawing a card stating the group they were in. In this article the two experimental groups are reported together as the aim was to assess the effect of exercise on RA patients. Nineteen subjects from the EG and eight from the CG completed the study.

### Intervention

The EG was required to train three times per week, 45 minutes at a time, for 12 weeks. The training was done under supervision of a biokineticist (specialised in physical training programmes). The CG received no intervention and was asked to continue with their sedentary lifestyles. The programme consisted of a warm-up phase (5 min), aerobic exercise (walking or aquatics for 20 min), strength training (10 min) and flexibility training (10 min). Aerobic exercise intensity was started at 60% of the heart rate maximum and was individually tailored to a maximum of 80%. Strength training in week 1 was 50% of the one repetition maximum and this was gradually increased to 80% in week 12.

### Measurements

Participants were tested at baseline and after 12 weeks (post-intervention). Measurements included:

- Physical parameters: height (cm); weight (kg); body mass index (BMI)
- Flexibility parameters: wrist flexion and extension (degrees); knee flexion and extension (degrees); hip flexion and extension (degrees); lateral flexion (side-ways bend [cm]); chair sit and reach (cm); scratch test (cm)
- Strength parameters: grip strength (kg); leg strength (kg); arm curls (s); sit to stand test (s)

- Fitness parameters: Rockport 1 mile walk test (min); VO<sub>2</sub>max Relative (ml/kg/min)
- Quality of life measures: Health Assessment Questionnaire (HAQ);<sup>18</sup> Visual Analogue Scale for pain (VAS)<sup>19</sup>
- Disease activity scores: DAS28(4) CRP version<sup>20</sup>

## Results

Collected data were captured and analysed by using SPSS Statistics 17.0. The non-parametric Mann-Whitney U test was used to compare the EG with the CG before and after intervention. The difference between the post-intervention and baseline measurements were used to assess whether the intervention had an effect. This method was used to control for initial differences between the two groups at baseline. To compare baseline measurements and post-intervention measurements within each group the Wilcoxon signed-rank test was used. Significant differences at the 5% and 10% level are reported.

Baseline characteristics for the two groups are shown in Table II.

A description of the measurements that changed significantly, as well as the direction that indicates an improvement, are explicated in Table III.

Firstly the functional parameters (flexibility, strength and fitness) will be reported on, whereafter the quality of life (HAQ and VAS) and disease activity parameters (DAS) will follow. Mean values are displayed in the tables. The graphs display the 95% confidence intervals around the mean.

### Functional parameters – between groups

At baseline the CG and EG were comparable, except for knee flexion. The EG had significantly better knee flexion, cf. Table IV.

### The training was done under supervision of a biokineticist (specialised in physical training programmes)

**Table I: Criteria for inclusion and exclusion**

Inclusion criteria	Exclusion criteria
Confirmed RA (ACR criteria) <sup>27</sup>	Smoking
Female gender	Diabetic disease
Controlled RA (stable medication for at least 3 months)	Systemic disease (cardiovascular, pulmonary, neurological, hepatic, kidney)
Medically fit to participate in an exercise programme	Drugs interfering with - autonomic nervous system (ANS) - cardiovascular system (CVS)
Willingness to participate in the study	Prior physical training programme
Age 30-60 years	Allergies to pool chemicals

**Table II: Baseline characteristics for the CG and the EG**

	CG (n=8)	EG (n=19)
Age, yrs	49.7 ± 4.3	47.3 ± 9.2
BMI	31.7 ± 4.7	25.3 ± 3.5
Medication use:		
Methotrexate	7	16
Sulfasalazine	0	2
Chloroquine	1	4
Leflunomide	0	2
Biologic drug	2	2
NSAIDs*	7	16
Prednisone	3	5


\*Non-steroidal anti-inflammatory drugs

**Table III: Parameter measurement descriptions**

Parameter	Descriptor	Direction of improvement
<b>Flexibility:</b>		
Knee flexion and extension	Degrees	↑
Hip flexion	Degrees	↑
Hip extension	Degrees	↓
Lateral flexion (sideways bend)	Middle finger distance from floor (cm)	↓
Chair sit and reach	Middle finger distance from toe (cm)	↑
Scratch test	Reaching behind back: distance from middle finger to inferior angle of scapula (cm)	↑
<b>Strength:</b>		
Hand grip	Dynamometer (kg)	↑
Leg strength	Dynamometer (kg)	↑
Arm curls	Time to execute 30 (s)	↓
Sit to stand	Time to rise from chair to standing position 20 times (s)	↓
<b>Fitness:</b>		
1 mile walk test	Time to walk 1 mile (min)	↓
VO <sub>2</sub> max	ml/kg/min	↑
<b>Quality of life:</b>		
VAS	Scale of 1-10	↓
HAQ	Scale of 1-3	↓
<b>Disease activity:</b>		
DAS28(4) CRP	Formula calculation	↓

**Table IV: Functional parameters – between groups**

	CG	EG	Group favoured
<b>At baseline</b>			
Right knee flexion (p-value = 0.003)	Mean = 124.25°	Mean = 138.26°	EG
Left knee flexion (p-value = 0.011)	Mean = 129°	Mean = 138.74°	EG
<b>Post-intervention (based on differences between post-intervention and baseline measurements)</b>			
Right lateral flexion (p-value = 0.014)	Mean = 2.88 cm (53.23-50.35)	Mean = 0.62 cm (47.63-47.01)	EG
Left lateral flexion (p-value = 0.015)	Mean = 1.84 cm (53.80-51.96)	Mean = -1.06 cm (47.54-48.60)	EG
Left chair sit and reach (p-value = 0.058)	Mean = 2.06 cm (-3.33 - (-)5.39)	Mean = 10.84 cm (3.56- (-)7.28)	EG
Right handgrip strength (p-value = 0.053)	Mean = 5.29 kg (24.25-18.96)	Mean = 7.76 kg (24.76-17.00)	EG
1 mile walk test (p-value = 0.011)	Mean = -0.90 min (17.57-18.47)	Mean = -1.98 min (14.93-16.91)	EG



**Graph 1: Functional parameters – post-intervention between groups**

Comparing the two groups post-intervention, right lateral flexion deteriorated for both groups, but significantly less so for the EG. Significant improvements were found in favour of the EG for left lateral flexion, chair sit and reach left, handgrip strength right and the 1 mile walk test (Table IV and Graph 1).

**Functional parameters – within groups**

The CG showed significant improvements for knee extension; however, hip extension and lateral flexion declined significantly from baseline to end of study. (Table Va and Graph 2a).

The strength parameters, including hand grip, leg strength, arm curls and sit to stand test also improved significantly from baseline to study completion in the CG (Table Vb and Graph 2b).

Within the EG the following flexibility parameters improved significantly post-intervention: knee flexion left, knee extension, hip flexion right, chair sit and reach and scratch test; however, as for the CG, hip extension declined significantly (Table VIa and Graph 3a).

The strength parameters that improved were hand grip, leg strength, arm curl test and sit to stand test (Table VIb and Graph 3b).

Both aerobic parameters i.e. the 1 mile walk test and the VO<sub>2</sub>maxRelative, also improved significantly from baseline to end of study in the EG (Table VIc and Graph 3c).

**Disease activity parameters – between groups**

The CG and EG did not differ at baseline with regard to their VAS, HAQ and DAS28(4) measurements. At completion of the study however, the EG improved significantly more than the CG with respect to DAS scores, while the CG rated themselves significantly better than the EG group did according to the HAQ (Table VII and Graph 4).

**Disease activity parameters – within groups**

The CG had significant improvements in the HAQ and VAS. The EG improved significantly in the DAS28(4) and VAS (Table VIII and Graph 5).


**Within the EG the flexibility parameters improved significantly post-intervention**

**Table Va: Functional parameters – flexibility variables within CG**


	Baseline	End of study	Difference	Outcome
Right knee extension (p-value = 0.05)	Mean = -4.13°	Mean = -1.63°	2.50°	Improved
Left knee extension (p-value = 0.013)	Mean = -5.63°	Mean = -2.13°	3.50°	Improved
Right hip extension (p-value = 0.046)	Mean = -21.25°	Mean = -17.88°	3.38°	Deteriorated
Right lateral flexion (p-value = 0.009)	Mean = 50.35 cm	Mean = 53.23 cm	2.88 cm	Deteriorated
Left lateral flexion (p-value = 0.018)	Mean = 51.96 cm	Mean = 53.80 cm	1.84 cm	Deteriorated

**Table Vb: Functional parameters – strength variables within CG**

	Baseline	End of study	Difference	Outcome
Right hand grip (p-value = 0.006)	Mean = 18.96 kg	Mean = 24.25 kg	5.29 kg	Improved
Left hand grip (p-value = 0.009)	Mean = 16.93 kg	Mean = 22.59 kg	5.66 kg	Improved
Leg strength (p-value = 0.04)	Mean = 58.49 kg	Mean = 68.19 kg	9.7 kg	Improved
Arm curls (p-value = 0.006)	Mean = 53.26 s	Mean = 33.17 s	-20.08 s	Improved
Sit to stand test (p-value = 0.009)	Mean = 64.29 s	Mean = 41.92 s	-22.37 s	Improved



Graph 2a: Functional parameters – within groups for CG: flexibility variables



Graph 2b: Functional parameters – within groups for CG: strength variables

Table VIa: Functional parameters - flexibility variables *within EG*


	Baseline	End of study	Difference	Outcome
Left knee flexion (p-value = 0.026)	Mean = 138.74°	Mean = 142.26°	3.53°	Improved
Right knee extension (p-value = 0.011)	Mean = -2.74°	Mean = -0.63°	2.11°	Improved
Left knee extension (p-value = 0.009)	Mean = -3.42°	Mean = -1.37°	2.05°	Improved
Right hip flexion (p-value = 0.06)	Mean = 71.47°	Mean = 75.11°	3.63°	Improved
Right hip extension (p-value = 0.06)	Mean = -12.95°	Mean = -8.37°	4.58°	Deteriorated
Left hip extension (p-value = 0.04)	Mean = -11.21°	Mean = -5.84°	5.37°	Deteriorated
Right chair sit and reach (p-value = 0.011)	Mean = -6.38 cm	Mean = 0.37 cm	6.76 cm	Improved
Left chair sit and reach (p-value < 0.001)	Mean = -7.27 cm	Mean = 3.56 cm	10.83 cm	Improved
Right scratch test (p-value = 0.007)	Mean = -4.76 cm	Mean = -0.25 cm	4.51 cm	Improved
Left scratch test (p-value = 0.01)	Mean = -8.48 cm	Mean = -4.97 cm	3.51 cm	Improved

Table VIb: Functional parameters - strength variables *within EG*

	Baseline	End of study	Difference	Outcome
Right hand grip (p-value < 0.001)	Mean = 16.99 kg	Mean = 24.76 kg	7.76 kg	Improved
Left hand grip (p-value < 0.001)	Mean = 16.56 kg	Mean = 22.53 kg	5.97 kg	Improved
Leg strength (p-value = 0.003)	Mean = 49.42 kg	Mean = 61.63 kg	12.21 kg	Improved
Arm curls (p-value < 0.001)	Mean = 51.54 s	Mean = 31.06 s	-20.48 s	Improved
Sit to stand test (p-value < 0.001)	Mean = 65.37 s	Mean = 36.88 s	-28.49 s	Improved


Table VIc: Functional parameters - aerobic variables *within EG*

	Baseline	End of study	Difference	Outcome
1 mile walk test (p-value < 0.001)	Mean = 16.90 min	Mean = 14.93 min	-1.97 min	Improved
VO <sub>2</sub> max Relative (p-value = 0.01)	Mean = 29.02 ml/kg/min	Mean = 30.83 ml/kg/min	1.80 ml/kg/min	Improved



**Graph 3a: Functional parameters – within groups for EG: flexibility variables\***  
 \* Left knee flexion and right hip flexion have been scaled by a factor 10 to facilitate the clustered plot on one set of axes

Research has shown that regular and controlled exercise for those whose disease is under control decreases joint pain and stiffness and improves joint mobility, strength and aerobic capacity




**Graph 3b: Functional parameters – within groups for EG: strength variables**

## Discussion


The results of this study, with a compliance rate of 77%, indicated significant improvements for flexibility, strength and aerobic capacity measured after a 12-week aerobic exercise intervention. Previous research indicates that exercise compliance for RA patients are on average between 50% and 95% depending on accessibility of the exercise, intensity, duration, cost, and comfort involved for the patient.<sup>21</sup> RA is the second-most common joint disease, causing various physical impairments either as result of the disease or due to inactivity.<sup>5,8</sup> Patients with RA are hesitant to get involved in any form of exercise because of fear of pain and disability. These fears are unfounded as research has shown that regular and controlled exercise for those whose disease is under control decreases joint pain and stiffness and improves joint mobility, strength and aerobic capacity without exacerbating pain or disease activity in persons with RA.<sup>3,21,22</sup>

The flexibility findings of this study support the findings of previous studies done on patients with RA.<sup>13,23,24</sup> The results showed significant improvements in flexibility of all the major joints for the EG whereas the flexibility of the CG remained relatively unchanged for the duration of the study.


Strength improved for both the CG and EG, even though the control group did not participate in a training programme. This is not an uncommon occurrence as a study by Bykerk and Keystone, showed similar results.<sup>25</sup> According to Häkkinen, the improvements in the CG can be explained by the learning effects of repeated physical testing and variation in symptoms.<sup>10</sup>



**Graph 3c: Functional parameters – within groups for EG: aerobic variables**



Graph 4: Disease activity parameters – post-intervention *between* groups



Graph 5: Disease activity parameters – *within* groups\*

\* VAS has been scaled by a factor 10 to facilitate the clustered plot on one set of axes

Table VII: Disease activity parameters – *between* groups\*

	CG	EG	Group favoured
DAS	Mean = -0.25 (p-value = 0.072) (3.1-3.35)	Mean = -0.78 (2.51-3.29)	EG
HAQ	Mean = -0.20 (p-value = 0.082) (0.36-0.56)	Mean = -0.16 (0.32-0.48)	CG

\* Differences between post-intervention and baseline measurements

The EG's strength improved by an average of 33% compared to the 27% improvement in the CG, indicating that the exercise intervention did have a positive effect on the patients' strength

Table VIII: Disease activity parameters – *within* groups

	Baseline	End of study	Difference	Outcome
<b>CG</b>				
HAQ (p-value = 0.032)	Mean = 0.56	Mean = 0.36	-0.20	Improved
VAS (p-value = 0.029)	Mean = 44.38	Mean = 31	-13.38	Improved
<b>EG</b>				
DAS (p-value < 0.001)	Mean = 3.29	Mean = 2.51	-0.78	Improved
VAS (p-value = 0.031)	Mean = 46.05	Mean = 35.58	-10.47	Improved

The EG's strength improved by an average of 33% compared to the 27% improvement in the CG, indicating that the exercise intervention did have a positive effect on the patients' strength. The average improvements in the EG for hand grip strength, leg strength, arm curl test and sit to stand test respectively were 29%, 20%, 40% and 44%. These results are in agreement with studies done by Van den Ende *et al.*, Häkkinen *et al.* and Stenstrom that similarly showed improvements in muscle strength ranging between 16% and 35%.<sup>13,24,26</sup>

Exercise programmes with the specific purpose of improving aerobic fitness have attracted only minimal attention in this population. Previous studies on the effect of aerobic exercise in RA patients made use of cycling, aquatics or aerobic dance.<sup>22,24,26</sup> Studies done by Iversen *et al*, Minor *et al*, and Van den Ende *et al* showed significant improvements in aerobic capacity (20%) following 12 weeks of dynamic exercise of medium to high intensities.<sup>6,23,24</sup> In keeping with these studies our results showed an average improvement of 14% in VO<sub>2</sub>max and 12% in

1 mile walk time in the EG, confirming that the intervention was successful in improving aerobic fitness. In contrast to these improvements, the CG showed a decrease in aerobic capacity of 30% over the same time period.

Results on the activities of daily living (HAQ) are difficult to explain, but are similar to previous publications. The improvement of disease activity is also in support of some previous studies.<sup>11-13,15,16</sup> Although the EG had marked improvement in their fitness parameters and disease activity scores, they did not improve significantly in their own assessment of activities of daily living. This might be because the inclusion criteria stipulated controlled disease and stable medication when the study started (HAQ scores were already low).

With improvements in flexibility, strength and aerobic fitness it can thus be concluded that the EG's functional capacity and disease activity improved as a result of a relatively short but well-controlled endurance exercise programme of 12 weeks. In agreement with Christie *et al.* exercise therapy should be considered the cornerstone of the multidisciplinary treatment approach of stable grade 1 or 2 RA.<sup>3</sup>

Although this study demonstrated that endurance exercise improves function and disease activity in RA patients, future research on the effect of exercise in female RA patients should aim for a larger study sample and a more individualised exercise programme in an attempt to decrease the SD between and within the groups for the tested parameters. Additionally the duration of intervention could be increased as very few studies lasted longer than 12 weeks thus the long-term effects of exercise on RA patients are not well known.<sup>22</sup> The long-term influence of an exercise programme after cessation of the intervention will also be informative.

*No benefits of any form have been received from a commercial party related directly or indirectly to the subject of this article.*

### Acknowledgements

I wish to acknowledge the biokineticists Natania Fourie and Anneke de Beer for their involvement with the training, Brenda Weder for the typing of the document and Maureen Brassel for helping with the literature search.

### References

- O'Dell JR. Rheumatoid arthritis: The clinical picture. In: Koopman WJ, Moreland LW (eds). *Arthritis and Allied Conditions: Textbook of Rheumatology*. 15<sup>th</sup> ed. Philadelphia: Lippincott, Williams & Wilkins 2005;1165-94.
- Armstrong L, Balady GJ, Berry MJ. *ACSM Guidelines for Exercise Testing and Prescription*. 7<sup>th</sup> ed. Philadelphia, Pennsylvania: Lippincott, Williams and Wilkins 2006;105-7.
- Christie A, Jamtvedt G, Dahm KT, Moe RH, Haavardsholm EA, Hagen KB. Effectiveness of nonpharmacological and nonsurgical interventions for patients with rheumatoid arthritis: an overview of systematic reviews. *Phys Ther* 2007 Dec 87;12:1697-715.
- Gaudin P, Leguen-Guegan S, Allenet B, Baillet A, Grange L, Juvin R. Is dynamic exercise beneficial in patients with rheumatoid arthritis? *Joint Bone Spine* 2008 Jan;75(1):11-17.
- Hicks JE. Rehabilitation strategies for patients with rheumatoid arthritis. *J Musculoskelet Med* 2000;17(4):991.
- Iversen MD. Physical therapy for management of RA: rehabilitation measures are most effective when started early. *J Musculoskelet Med* 2002;19(9):352-58.
- Giannini MJ, Protas EJ. Exercise response in children with and without juvenile rheumatoid arthritis: a case-comparison study. *Phys Ther* 1992 May;72(5):365-72.
- Millar AL. Action plan for arthritis: Your guide to pain-free movement. *Human Kinetics* 2003;31-49.
- Gossec L, Pavy S, Pham T, Constantin A, Poiraudou S, Combe B, *et al.* *Nonpharmacological treatments in early rheumatoid arthritis: clinical practice guidelines based on published evidence and expert opinion*. 2006 Jul;73(4):396-402.
- Häkkinen A. Effectiveness and safety of strength training in rheumatoid arthritis. *Curr Opin Rheumatol* 2004 Mar;16(2):132-7.
- Van Den Ende CH, Vliet Vlieland TP, Munneke M, Hazes JM. Dynamic exercise therapy for rheumatoid arthritis. *Br J Rheumatol* 1998;37(6):677-87.
- Flint-Wagner HG, Lisse J, Lohman TG, Going SB, Guido T, Cussler E, *et al.* Assessment of a sixteen-week training program on strength, pain, and function in rheumatoid arthritis patients. *J Clin Rheumatol* 2009 Jun;15(4):165-71.
- Stenstrom CH. Home exercise in rheumatoid arthritis functional class II: goal setting versus pain attention. *J Rheumatol* 1994 Apr;21(4):627-34.
- Lee EO, Kim JI, Davis AH, Kim I. Effects of regular exercise on pain, fatigue, and disability in patients with rheumatoid arthritis. *Fam Community Health* 2006 Oct-Dec;29(4):320-7.
- Lynberg K, Danneskiold-Samsøe B, Halskov O. The effect of physical training on patients with rheumatoid arthritis: changes in disease activity, muscle strength and aerobic capacity. A clinically controlled minimized cross-over study. *Clin Exp Rheumatol* 1988 Jul-Sep;6(3):253-60.
- Neuberger GB, Aaronson LS, Gajewski B, Embretson SE, Cagle PE, Loudon JK, *et al.* Predictors of exercise and effects of exercise on symptoms, function, aerobic fitness, and disease outcomes of rheumatoid arthritis. *Arthritis Rheum* 2007 Aug 15;57(6):943-52.
- Häkkinen A, Häkkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994;23(5):237-42.

18. Pincus T, Summey JA, Soraci SA Jr, Wallston KA, Hummon NP. Assessment of patient satisfaction in activities of daily living using a modified Stanford Health Assessment Questionnaire. *Arthritis Rheum* 1983 Nov;**26**(11):1346-53.
19. Callahan LF, Pincus T. A clue from a self-report questionnaire to distinguish rheumatoid arthritis from noninflammatory diffuse musculoskeletal pain. The P-VAS:D-ADL ratio. *Arthritis Rheum* 1990 Sep;**33**(9):1317-22.
20. Prevoe ML, van't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, van Riel PL. Modified disease activity scores that include twenty-eight joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995 Jan;**38**(1):44-8.
21. Westby MD, Li L. Physical therapy and exercise for arthritis: Do they work? *Geriatrics Aging* 2006;**9**(9):624-30.
22. Metsios GS, Stavropoulos-Kalinoglou A, Veldhuijzen van Zanten JJ, Treharne GJ, Panoulas VF, Douglas KM, *et al.* Rheumatoid arthritis, cardiovascular disease and physical exercise: a systematic review. *Rheumatology* (Oxford) 2008 Mar;**47**(3):239-48.
23. Minor MA, Hewett JE, Webel RR, Anderson SK, Kay DR. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum*. 1989 Nov;**32**(11):1396-405.
24. van den Ende CH, Hazes JM, le Cessie S, Mulder WJ, Belfor DG, Breedveld FC, *et al.* Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomised clinical trial. *Ann Rheum Dis* 1996 Nov;**55**(11):798-805.
25. Bykerk VP, Keystone EC. What are the goals and principles of management in the early treatment of rheumatoid arthritis? *Best Pract Res Clin Rheumatol* 2005 Feb;**19**(1):147-161.
26. Häkkinen A, Sokka T, Kotaniemi A, Kautiainen H, Jappinen I, Laitinen L, *et al.* Dynamic strength training in patients with early rheumatoid arthritis increases muscle strength but not bone mineral density. *J Rheumatol* 1999 Jun;**26**(6):1257-1263.
27. Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, *et al.* The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988 Mar;**31**(3):315-324

• SAOJ

---

## Criteria for authorship of articles

---

The following are internationally acknowledged criteria for authors/co-authors.

With the increase in faculty and in research projects, there is a potential for increased confusion and conflict regarding appropriate authorship credit on manuscripts and presentations. The following are some relatively standardised criteria that can be helpful. These may be overstrict when considering clinical studies in which surgeons often do the "hands on work" that create the study but may not perform major analysis and writing functions. However, all authors should read and contribute editing comments prior to submission.

### Relman criteria for authorship

In particular, to qualify as an author a person should fulfil at least three of the following five requirements:

1. Conception of idea and design of experiment
2. Actual execution of experiment; hands on lab work
3. Analysis and interpretation of data
4. Actual writing of manuscript
5. Be able to present to a learned gathering a lecture on the work; interpret it, defend it and take responsibility for it.

These are just guidelines. On the other hand it is probably far worse to leave someone off the list who feels they may have contributed than to include someone who did a bit less.

We should all be as inclusive as possible, offer our interested colleagues opportunity to provide input, analysis and editing of our works to support each other and make our papers better.

---