

Effect of Moistube and subsurface drip irrigation on cowpea (*Vigna unguiculata* (L.) Walp) production in South Africa

Edwin Kimutai Kanda^{1,3}, Aidan Senzanje¹ and Tafadzwanashe Mabhaudhi²

¹School of Engineering, University of KwaZulu-Natal, P. Bag X01, Scottsville, Pietermaritzburg, South Africa

²Centre for Transformative Agricultural and Food Systems, School of Agricultural, Earth and Environmental Sciences, University of KwaZulu-Natal, P. Bag X01, Scottsville, 3209, Pietermaritzburg, South Africa

³Department of Civil and Structural Engineering, Masinde Muliro University of Science and Technology, PO Box 190, Kakamega, 50100, Kenya

Moistube irrigation (MTI) is a new subsurface irrigation technology where the water emits from a semi-permeable membrane at a slow rate depending on applied pressure and soil water potential. There is lack of information on how various crops respond to MTI. This study determined growth, yield and water use efficiency (WUE) of cowpea (*Vigna unguiculata* (L.) Walp) under varying water regimes under MTI and subsurface drip irrigation (SDI), using field and glasshouse experiments in summer and winter of 2018, respectively. A split-plot design arranged in randomized complete blocks, replicated 3 times, with SDI as the control experiment was used. The main plot was irrigation type while the sub-plots were the water regimes. The water treatments consisted of full irrigation (100% of crop water requirement (ET_c)), and deficit irrigation (DI) of 70% ET_c and 40% ET_c. Water deficit had a significant effect ($p < 0.05$) on time to flowering; plants under 40% ET_c flowered 14 days earlier than plants at 100% ET_c. There were significant ($p < 0.05$) differences in yield components. Grain yields were 1 280 kg·ha⁻¹, 2 401 kg·ha⁻¹ and 3 189 kg·ha⁻¹ for 40% ET_c, 70% ET_c and 100% ET_c, respectively, but no significant ($p > 0.05$) differences were recorded between SDI and MTI. However, at 40% ET_c, SDI had 15% higher yield than MTI. Biomass varied significantly ($p < 0.05$) with irrigation type and water treatment. Grain WUE varied significantly ($p < 0.05$) among the water regimes. The highest WUE was achieved under SDI at 70% ET_c but was not significantly different from that under MTI at 70% ET_c. In conclusion, performance of cowpea was similar under the two irrigation systems under moderate DI but was better for SDI under severe DI with respect to biomass and WUE for the summer trial. Moderate DI improved the grain WUE while all the DI conditions improved the biomass WUE.

CORRESPONDENCE

Edwin Kimutai Kanda

EMAIL

Kandaedwin@gmail.com

DATES

Received: 3 May 2019

Accepted: 3 April 2020

KEYWORDS

crop growth
deficit irrigation
indigenous legume
irrigation
semi-permeable membrane
water use efficiency
yield

COPYRIGHT

© The Author(s)

Published under a Creative

Commons Attribution 4.0

International Licence (CC BY 4.0)

INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp) is one of the most important legumes grown in most parts of the world (Sebetha et al., 2010). It is the most commonly cultivated crop in resource-scarce countries of Africa, Asia, Central and South America, due to its ability to withstand extremely harsh environmental conditions such as high temperatures, limited water availability and poor soil fertility (Shiringani and Shimelis, 2011). It is also grown in European countries around the Mediterranean, such as Turkey (Basaran et al., 2011, Peksen, 2007). In South Africa, cowpea is cultivated in Limpopo, KwaZulu-Natal, Mpumalanga and the North West Provinces (DAFF, 2014). It can also be found in the wild in KwaZulu-Natal, Mpumalanga and Limpopo Provinces (Van Rensburg et al., 2007). Cowpea is nutritionally valuable in humans and animals. It is consumed as grains (dry and fresh) and vegetable leaves (Badiane et al., 2004) whereas the haulms are utilized as forage for livestock (Sprent et al., 2009). Cowpea grains are rich in proteins which could complement the diets of the majority of African households whose diets mainly consist of starch (Singh et al., 2003). Therefore, enhanced production of this crop would help in alleviating food insecurity in Africa.

Most of cowpea production is under rain-fed systems by small-scale farmers (Singh et al., 2003). Unavailability of rainfall or non-uniform distribution thereof means that yields cannot be guaranteed since water deficits affect plant growth and flowering (Timko and Singh, 2008). Indeed, studies have demonstrated that water deficit at flowering negatively affected yields of cowpea (Abdoul Karim et al., 2018; Ahmed and Suliman, 2010; Anyia and Herzog, 2004; Peksen, 2007). Households depending on cowpea are consequently exposed to risks of crop failure, hunger and malnutrition. Irrigation helps in stabilizing yields and acts as insurance to farmers in instances where there is rainfall variability or insufficient rainfall to meet crop requirements. Irrigation also allows for all-year round production, especially in the tropics and subtropics where temperatures are favourable for cowpea growth.

However, irrigation is the biggest consumer of freshwater resources, accounting for about 70% of total water use in arid and semi-arid areas (Fereser and Soriano, 2007). The irrigation sector in South Africa contributes about 60% of total water use (Reinders et al., 2010). This, therefore, requires adoption of efficient irrigation systems and appropriate agricultural water management practices to save water. Deficit irrigation (DI) is one of the main water-saving irrigation strategies where the

volume of water applied is below the crop water requirement, with the aim of maximizing crop water productivity (Feres and Soriano, 2007). Adoption of efficient irrigation methods such as subsurface drip irrigation (SDI) helps in reducing agricultural water use by minimizing non-beneficial components such as soil evaporation, runoff and drainage (Ayars et al., 1999).

Moistube irrigation (MTI) is a relatively new type of irrigation technology which originated in China. It is similar to SDI but, instead of flowing from emitters, water flows out of the Moistube membrane as a function of applied pressure and the soil water potential (Yang et al., 2008). Some studies have shown that MTI saves water and has higher water use efficiency (WUE) than conventional irrigation methods. For instance, a comparison between MTI and SDI for tomatoes showed that the former resulted in similar yield per unit area compared to the latter but also improved WUE by 13% (Xue et al., 2013). Lyu et al. (2016) found that MTI can achieve about 38% water savings compared to drip irrigation with mulch in production of tomatoes. Besides an increase in total yield, tomato quality in terms of fruit diameter, weight, Vitamin C, soluble sugar and soluble acid ratio were 8.6%, 12%, 27%, 4.5% and 21% higher, respectively, using MTI compared to drip irrigation. Yao et al. (2014) compared conventional irrigation, MTI, and rainfed water conditions. In the study, Moistube-irrigated navel oranges achieved the highest leaf respiration index, photosynthetic rate, specific leaf area and quantum yield. A study by Zhang et al. (2016a) established that water savings could be achieved in cabbage production using MTI, but the results could not be verified and thus further studies were recommended. Yin et al. (2017) reported 21% water savings in MTI-irrigated spinach compared to conventional irrigation. However, Zhang et al. (2017) found that MTI decreased maize yield significantly relative to that obtained under SDI, while wheat yield only tended to decrease. In the same study, there were no significant differences in crop WUE between MTI and SDI.

Based on the above information, and the potential importance of cowpea in South Africa, it was considered necessary to evaluate the response of cowpea under the new water-saving irrigation technology (MTI) and conventional irrigation (SDI). The aim of this study, therefore, was to determine the growth, development, yield and water use efficiency of cowpea under varying water regimes under MTI and SDI. Correlation analysis was done to determine the relationships among the growth and yield variables. This study was based on two hypotheses. First, it was hypothesized that there was no significant difference between the response of cowpea under MTI and SDI, and secondly, that WUE of cowpea could be improved by a DI strategy.

MATERIALS AND METHODS

Glasshouse experiment (summer season)

This research was carried out in tunnels at the Controlled Environment Facility (CEF) of UKZN, Pietermaritzburg Campus (29.58° S, 30.42° E) during summer 2018. The experiment was carried out in a 11 m long glasshouse in raised beds measuring 0.75 m wide and 0.75 m high. The soil texture was loam (42.3% sand, 33.3% silt, 24.4% clay) with a bulk density of 1.36 g·cm⁻³.

The experiment was laid out in a split-plot design arranged in randomized complete blocks. The main block was the irrigation type (SDI and MTI) while the sub-plots were 3 water regimes replicated 3 times. The water regimes imposed consisted of irrigation to meet the full crop water requirement (100% ET_c), and DI of 70% ET_c and 40% ET_c. The drip emitters and Moistube tapes were installed at a 15 cm depth, which was guided by the crop rooting depth and the need to have emitters close to the surface for crop establishment.

Cowpea (brown mix variety) was planted on 14 February 2018 (summer season). The plant spacing was 50 cm between rows and 30 cm within rows giving a plant density of 66 667 plants·ha⁻¹. A soil fertility test conducted at Cedara Agricultural College indicated that the soil required phosphorus, which was applied as Single Superphosphate (10.5% P) at 60 kg·ha⁻¹. The DI was induced from 21 days after planting (DAP).

Other agronomic management practices such as weed, pest and disease control were done according to recommended best practices developed by the then Department of Agriculture, Forestry and Fisheries of the Republic of South Africa (DAFF, 2014).

Field experiment (winter season)

Field trials were conducted at the University of KwaZulu-Natal's Ukulinga Research Farm in Pietermaritzburg (29.67° S, 30.41° E) during winter 2018. These were carried out in a 12 m by 5 m tunnel where the soil texture was clay (24.3% sand, 23.6% silt and 52.1% clay) with bulk density of 1.23 g·cm⁻³. The tunnel at Ukulinga had open ends with free movement of air to imitate field conditions as much as possible. The temperature in the tunnels varied from 4°C to 15°C during the growing period.

The experimental layout, cowpea variety, plant spacing, Moistube and SDI placement depth were similar to that described for the glasshouse experiment. The crop was planted on 25 May 2018 (winter season). Soil fertility test indicated that the soil did not have a nutrient deficiency and, therefore, fertilizers were not added. The DI water regimes were introduced 30 DAP when the crops were fully established. The long crop establishment period was due to the low temperatures which affected emergence. Other agronomic management practices were done according to recommended best practices as described in the glasshouse experiment.

Estimation of crop water requirements

The crop water requirements (ET_c) for each crop growth stage were determined using potential reference evapotranspiration and crop coefficients as described in Allen et al. (1998). The crop coefficients adopted for cowpea were 0.4, 1.1 and 0.4 for the initial, mid- and end-growth stage, respectively.

The net irrigation requirement (I_{net}) was calculated as ET_c less rainfall. Since rainfall equalled zero, the I_{net} was equal to ET_c. The water use efficiency was determined as the ratio of the yield to the amount of irrigation applied.

The different water regimes were applied by varying the irrigation interval in such a way that the total amount of irrigation was 100%, 70% and 40% of ET_c. In SDI, the amount of water applied per irrigation event was the same, but the irrigation interval was different for the DI, i.e., 100% ET_c < 70% ET_c < 40% ET_c. A drip emitter of nominal flow rate of 1.6 L·h⁻¹ was used in this study. This flowrate was used to calculate the amount of water to be applied at every irrigation event. The flow from Moistube was in the range of 0.24 L·h⁻¹·m⁻¹ at 20 kPa to 1.73 L·h⁻¹·m⁻¹ at 100 kPa (Kanda et al., 2018). The pressure was adjusted according to the crop water requirement. MTI was supposed to be continuous, i.e., water applied throughout the cropping cycle but the pressure regulators available were not sufficiently low to allow for continuous water application. Therefore, the water application was applied intermittently ranging from 3 days continuously per dekad, 5 days and 8 days per dekad for 40% ET_c, 70% ET_c and 100% ET_c, respectively.

Data collection and analysis

Weather data were obtained inside the tunnel using HOBO data logger sensors (Onset Computer Corporation, USA). The variables measured were temperature, relative humidity and solar radiation, while wind speed was measured using a Kestrel 3000 anemometer (Nielsen-Kellerman, Inc. USA) which were mounted 2 m above the ground.

Leaf area index (LAI) was measured weekly using the LAI2200 canopy analyser (LI-COR Inc. USA). The time to 50% flowering was determined by counting the number of flowered plants and was taken as days elapsed until 50% of the plants in each plot had flowered. Determination of yield components was done by sampling 10 plants per plot excluding border plants. All the pods from each plant were harvested, counted and then shelled for yield analysis. Aboveground biomass of each of the 10 plants per plot was determined by cutting each plant and then weighing after air drying for 3 weeks. For the field experiment, 5 plants in the inner row, excluding border plants, were harvested and weighed after air drying to determine the biomass.

The harvest index (HI) was computed using Eq. 1 (Cisse, 2001):

$$HI = \frac{\text{Grain yield (kg ha}^{-1}\text{)}}{\text{Biomass (kg ha}^{-1}\text{)}} \times 100\% \quad (1)$$

The water use efficiency (WUE) was computed using Eq. 2 (Albaji et al., 2011):

$$WUE = \frac{\text{Yield or Biomass (kg ha}^{-1}\text{)}}{\text{Amount of irrigation applied (m}^3\text{ha}^{-1}\text{)}} \quad (2)$$

The amount of irrigation applied was the cumulative irrigation depth (mm) applied during the irrigation events and then converted to volume per hectare.

The measured data was analysed using GenStat version 18 (VSN International, Hemel Hempstead, UK) to determine ANOVA of the variables. Separation of means of significant variables was done using Duncan's Least Significant Differences (LSD) at a 5% significance level in GenStat. Correlation analyses were carried out on growth and yield components to determine the relationship between variables.

RESULTS AND DISCUSSION

Leaf area index

LAI is an important growth parameter as it signifies the extent of the assimilative capacity of a crop under existing

environmental conditions (Farooq et al., 2012). The LAI varied among the treatments in both experiments. In the summer trials, there were no significant differences between the two types of irrigation (Table 1), but LAI under SDI at 40% ET_c was significantly ($p < 0.05$) higher than MTI at 40% ET_c (Fig. 1). There was no significant difference between SDI and MTI at full irrigation (100% ET_c) and moderate (70% ET_c) deficit irrigation ($p > 0.05$). The LAI was significantly lower under MTI at 40% ET_c compared to both SDI and MTI at 100% ET_c water regimes (Table 1).

The LAI for the winter trials showed variations among treatments and irrigation type (Table 2). The mean seasonal LAI was not significantly ($p > 0.05$) different between MTI and SDI at 100% ET_c. However, LAI under MTI was significantly lower than SDI at 70% ET_c. At 40% ET_c the LAI became significantly lower compared to the wetter regimes after 10 weeks (Fig. 2). The LAI of 100% ET_c MTI plants was significantly ($p < 0.05$) lower than where 100% and 70% ET_c SDI was applied after 10 weeks (Fig. 2).

Full irrigation in the summer season experiments resulted in a seasonal peak LAI of 4 and 3.88 for MTI and SDI, respectively (Fig. 1). Applying MTI at 40% ET_c resulted in the lowest seasonal peak LAI. The seasonal mean LAI of MTI plants irrigated at 40% ET_c decreased significantly (25.6%) compared to those receiving 100% ET_c MTI and SDI ($p < 0.05$) (Table 1). These results were consistent with those reported by Souza et al. (2017) where LAI declined by between 13% and 47% due to water deficit. The reduction in LAI could be attributed to decreased leaf appearance rate and abscission which are considered as drought avoidance mechanisms (Abayomi and Abidoye, 2009). Reduction in leaf area due to water deficit arises because of inhibited cell growth (Fathi and Tari, 2016). According to Prasad et al. (2008), mild water deficit causes reduction in leaf number, retarded leaf expansion rate and reduced leaf size while severe water deficit inhibits leaf appearance.

Significant ($p < 0.05$) seasonal differences in cowpea growth between the summer and winter trials were reflected by the seasonal LAI trends. During winter, most of the growth parameters were affected by the night temperatures, which were in most cases very low ($< 10^\circ\text{C}$). The low temperatures significantly delayed time to emergence (≈ 15 days) and led to slower leaf area development ($\text{LAI} \leq 1$) compared to summer trials where cowpea emerged after an average of 7 days and reached a higher LAI (≥ 1.9). As illustrated in Figs 1 and 2, the maximum LAI was significantly lower ($p < 0.05$) for the winter compared to the summer experiment. Ntombela (2012) found that low temperature limited growth characteristics

Table 1. Effect of irrigation type and deficit irrigation on LAI for the summer experiment

Irrigation	Water regime	Mean seasonal LAI
Moistube	100% ET _c	2.59 ^a
SDI	100% ET _c	2.55 ^a
Moistube	70% ET _c	2.39 ^{ab}
SDI	70% ET _c	2.40 ^{ab}
Moistube	40% ET _c	1.93 ^b
SDI	40% ET _c	2.21 ^{ab}
LSD (irrigation)		0.3062
LSD (ET _c)		0.3751
LSD (irrigation x ET _c)		0.5304

Mean values in same column followed by same superscript letter do not significantly differ at 5% level of significance by LSD. Data in parentheses are the standard deviations

Table 2. Effect of irrigation type and deficit irrigation on LAI for the winter experiment

Irrigation	Water regime	Mean seasonal LAI
Moistube	100% ET _c	0.41 ^{abc}
SDI	100% ET _c	0.55 ^a
Moistube	70% ET _c	0.36 ^c
SDI	70% ET _c	0.51 ^{ab}
Moistube	40% ET _c	0.31 ^c
SDI	40% ET _c	0.37 ^{bc}
LSD (irrigation)		0.0795
LSD (ET _c)		0.0973
LSD (irrigation x ET _c)		0.1377

Mean values in same column followed by same superscript letter do not significantly differ at 5% level of significance by LSD. Data in parentheses are the standard deviations

of cowpea more than water deficit. Therefore, irrespective of water availability, temperature is a significant factor in cowpea growth. This may be the reason for its popularity in tropical and subtropical countries in Sub-Saharan Africa, Asia, central and South America (Singh et al., 2003).

Time to flowering

Water deficit affects crop growth and development by not only retarding cell division and expansion but also by altering the initiation and duration of phenological stages (Prasad et al., 2008). Time to flowering is an environmental adaptive feature of most annual crops (Ishiyaku et al., 2005). In the present study, the cowpea grown in winter failed to flower due to low temperatures, which was consistent with results reported by Ilunga (2014) where cowpea planted in June (winter season), at the same site, failed to flower.

In the summer experiment, the number of days to 50% flowering varied significantly ($p < 0.05$) across the water regimes, as shown in Table 3. Cowpea flowered earlier at 40% ET_c for both MTI and SDI compared to 70% ET_c for SDI and the fully irrigated treatments. Full irrigation under MTI significantly delayed time to 50% flowering (10 days) compared to plants subjected to severe deficit (40% ET_c) and MTI 70% ET_c . The latter treatment flowered 4 days later than MTI 40% ET_c , but they were not significantly

different ($p > 0.05$). Cowpea receiving SDI at 100% ET_c took significantly longer to reach the 50% flowering stage than plants subjected to all the other water regimes.

The accelerated time to flowering due to water deficit reported in this study was consistent with that reported by Ilunga (2014) for the same variety (mixed brown), where rain-fed induced water deficit enhanced time to flowering by 7 days. The MTI supplies water to the crop at 80–90% of field capacity (Zhang et al., 2012) and hence a subsequent decrease in water, e.g. 70% ET_c , results in slightly lower water availability to the crop than at 70% ET_c under SDI. This could possibly explain the non-significant difference between time to 50% flowering under 70% ET_c SDI and 100% ET_c under MTI. The shorter duration to flowering as a result of water deficit is considered to be a drought escape mechanism in cowpea (Ehlers and Hall, 1997). However, some studies have reported delayed flowering due to water stress (Abayomi and Abidoye, 2009; Faloye and Alatis, 2017; Ntombela, 2012). The response of time to flowering under water deficit in cowpea depends on the genotype, as found by Dadson et al. (2005) where some exhibited early flowering while others had delayed flowering.

Yield and yield components

The cowpea at the field trial failed to flower due to low temperatures during the winter season and, therefore, the results reported here for the yield components and biomass were for the summer experiment, while only biomass was reported for the winter season.

Irrigation system type and deficit irrigation levels affected yield and yield components of the summer trial significantly (Table 4). Irrigation system type did not affect pod number or mass significantly, nor seed mass per plant. It also had no significant effect on grain yield or shelling percentage. SDI compared to MTI tended to increase biomass per plant and per hectare, but these were only significantly higher at the 70% and 40% ET_c deficit irrigation levels.

Deficit irrigation at 70% and 40% ET_c decreased pods, pod mass, seed mass and biomass per plant as well as grain yield significantly, compared to where irrigation was applied at 100% ET_c , irrespective of irrigation system type. Compared to SDI at 100% ET_c , a significant reduction in yield (57.7%) was recorded at 40% ET_c under MTI while drip at 40% ET_c led to a decline in yield by 50.2%. Similarly, the decline in yield at 70% ET_c relative




Figure 1. Leaf area index for different irrigation treatments of the summer experiment




Figure 2. Leaf area index for different irrigation treatments of the winter experiment

Table 3. Effect of irrigation system and deficit irrigation on cowpea time to flowering

Irrigation type	Water regime	Time to flowering (days)
Moistube	100% ET_c	65.7 (3.06) ^b
SDI	100% ET_c	74.3 (4.16) ^c
Moistube	70% ET_c	59.7 (4.51) ^a
SDI	70% ET_c	67.3 (2.08) ^b
Moistube	40% ET_c	55.3 (2.52) ^a
SDI	40% ET_c	56.7 (2.08) ^a
LSD (Irrigation)		3.3
LSD (ET_c)		4.04
LSD (Irrigation x ET_c)		5.72
CV (%)		5.1

Mean values in the same column followed by the same superscript letter do not significantly differ at 5% level of significance by LSD. Data in parentheses are the standard deviations

to the fully irrigated crop was 20.5% and 13.9% under MTI and SDI, respectively.

The results were similar to those found by Mousa and Qurashi (2017), where water deficit reduced the number of pods per plant. Abayomi and Abidoye (2009) also found reduced pod weight and number of seeds per plant due to water deficit. In another study, Hamidou et al. (2007) reported an average reduction of 60% in the number of pods per plant due to water deficit. The reduction in the number of pods per plant could be attributed to a lower number of flower buds and loss of flowers due to water deficit in the reproductive stage (Abdoul Karim et al., 2018; Maleki et al., 2017).

Deficit irrigation reduced yields, mainly due to significantly lower LAI, pod number per plant, pod mass per plant and harvest index. Maleki et al. (2017) reported a decline of between 4% and 59% in cowpea yields under a water regime of 40–80% of full irrigation. Reduction in yield due to water deficit is attributed to reduced photosynthetic active radiation absorption rate by plants and reduction in radiation efficiency (Fathi and Tari, 2016). According to Prasad et al. (2008), water deficit generally reduces grain yields, since it affects biomass production prior to flowering and negatively affects the reproduction phase of pollination as well as biomass partitioning during yield formation. Water deficit leads to a decline in leaf expansion, lower production of leaves and leaf senescence, which ultimately decreases the biomass (Figueiredo et al., 2001). Besides reduced leaf area, Anyia and Herzog (2004) associated decline in biomass of cowpea with reduced leaf gas exchange due to water deficit.

The significantly low temperatures during winter reduced the biomass compared to the summer season (Tables 4 and 5). There was no significant difference in biomass between the two irrigation system types. However, there were significant differences among the three water regimes where 40% ET_c resulted in significantly lower biomass compared to 70% ET_c and 100% ET_c. However, biomass at 70% ET_c and 100% ET_c were not significantly different. Biomass was significantly correlated to LAI ($r = 0.76, p < 0.05$), as in Table 6. The low biomass produced during the winter season (Table 5) may therefore be attributed to low LAI (Fig. 2). The low LAI was contributed by cold stress which inhibited leaf expansion. The findings concurred with those found by Ntombela (2012) where cold temperatures significantly reduced cowpea yield compared to water deficit.

Yield is a product of several components such as number of germinated plants, dry matter partitioning, numbers of seeds, and size of the seeds (Prasad et al., 2008). There was a strong positive correlation between the total grain yield and number of pods per plant ($r = 0.97$), pod mass per plant ($r = 0.96$), number of seeds per plant ($r = 0.96$), biomass and harvest index ($r = 0.94$) as shown in Table 6. Among these attributes, pod mass and harvest index had a significant contribution to grain yield ($p < 0.05$). Total grain yield ($r = 0.85$) and biomass ($r = 0.76$) were significantly correlated with LAI ($p < 0.05$). However, yield was strongly and negatively correlated with the number of days to 50% flowering ($r = -0.80$).

Table 4. Effect of irrigation system type and deficit irrigation on yield and yield components for the summer trial

Irrigation	Water regime	Pods/ plant	Pod mass/ plant (g)	Seeds/ plant	Seed mass/ plant (g)	Biomass/ plant (g)	Grain yield (kg-ha ⁻¹)	Biomass (kg-ha ⁻¹)	Harvest Index (%)	Shelling (%)
Moistube	100% ET _c	24 (7) ^c	67.8 (16.7) ^c	350 (94) ^c	48.3 (9.6) ^c	139.1 (18.7) ^e	3 189 (634) ^c	9 272 (1 247) ^e	34.8 (5.3) ^c	72.4 (8.0) ^{ab}
SDI	100% ET _c	25 (7) ^c	67.3 (20.7) ^c	345 (101) ^c	45.8 (10.5) ^c	145.2 (16.5) ^e	3 025 (695) ^c	9 678 (1 098) ^e	31.5 (6.2) ^b	70.5 (11.7) ^{ab}
Moistube	70% ET _c	19 (6) ^b	50.1 (17.6) ^b	268 (95) ^b	36.4 (9.4) ^b	120.2 (15.7) ^c	2 401 (612) ^b	8 012 (1 048) ^c	30.1 (5.5) ^b	75.1 (13.0) ^b
SDI	70% ET _c	19 (6) ^b	52.1 (15.8) ^b	315 (96) ^c	39.5 (10.6) ^b	128.9 (20.1) ^d	2 605 (701) ^b	8 590 (1 339) ^d	30.5 (6.0) ^b	77.3 (11.8) ^b
Moistube	40% ET _c	11 (5) ^a	29.9 (19.4) ^a	162 (84) ^a	19.4 (9.1) ^a	85.5 (10.0) ^a	1 280 (598) ^a	5 701 (926) ^a	22.4 (9.2) ^a	71.23 (20.6) ^{ab}
SDI	40% ET _c	13 (5) ^a	36.7 (15.3) ^a	175 (60) ^a	22.8 (7.0) ^a	100.4 (18.9) ^b	1 505 (462) ^a	6 694 (1 263) ^b	22.6 (4.8) ^a	66.54 (17.8) ^a
LSD (Irrigation)		1.8	5.2	9.5	2.6	5.3	88.3	350.6	1.72	2.6
LSD (ET _c)		2.3	6.4	11.6	3.2	6.44	108.1	429.4	2.1	3.1
LSD (Irr x ET _c)		3.2	9.0	16.4	4.6	9.11	152.9	607.2	2.9	4.4
CV (%)		24.1	24.5	23.3	20.9	15.3	20.9	15.3	17.4	13.3

Mean values in the same column followed by the same superscript letter do not significantly differ at 5% level of significance by LSD. Data in parentheses are the standard deviations

Table 5. Effect of irrigation system type and deficit irrigation on biomass for the winter trial

Irrigation	Water regime	Biomass/plant (g)	Total biomass (kg-ha ⁻¹)
Moistube	100% ET _c	27.36 (8.97) ^b	1 824 (598.12) ^b
SDI	100% ET _c	31.11 (9.15) ^b	2 074 (610.03) ^b
Moistube	70% ET _c	26.10 (7.39) ^b	1 740 (492.38) ^b
SDI	70% ET _c	28.53 (8.99) ^b	1 902 (599.99) ^b
Moistube	40% ET _c	17.38 (7.82) ^a	1 159 (521.34) ^a
SDI	40% ET _c	10.94 (4.88) ^a	729 (325.3) ^a
LSD (Irrigation)		3.943	262.9
LSD (ET _c)		4.829	322.0
LSD (Irrigation x ET _c)		6.830	455.3
CV (%)		24.3	24.3

Mean values in the same column followed by the same superscript letter do not significantly differ at 5% level of significance by LSD. Data in parentheses are the standard deviations

Table 6. Correlation analysis of growth and yield components for CEF experiment

	Pod No.	Pod mass/ Plant	Seeds/ plant	Grain yield (kg·ha ⁻¹)	Biomass (kg·ha ⁻¹)	Harvest index (%)	LAI	Days to flowering
Pods	1.00							
Pod mass/plant	0.99	1.00						
Seeds/plant	0.96	0.97	1.00					
Grain yield (kg·ha ⁻¹)	0.97	0.96	0.96	1.00				
Biomass (kg·ha ⁻¹)	0.98	0.98	0.98	0.98	1.00			
Harvest index (%)	0.95	0.94	0.95	0.94	0.93	1.00		
LAI	0.85	0.85	0.78	0.85	0.76	0.89	1.00	
Days to flowering	-0.83	-0.81	-0.77	-0.80	-0.71	-0.78	-0.77	1.00

Water use efficiency

Grain WUE varied significantly among the water regimes for the summer crop (Table 7). The highest grain WUE was achieved under SDI at 70% ET_c, but it was comparable to that of plants receiving MTI at 70% ET_c. MTI at 40% had the lowest grain WUE. Irrigation type did not significantly affect grain WUE at 100% ET_c. With respect to SDI, DI of 70% ET_c and 40% ET_c improved grain WUE relative to the fully irrigated crop by 17.3% and 4.9%, respectively. In MTI, DI improved grain WUE by 4.1% at 70% ET_c, but it decreased by 17% at 40% ET_c. This shows that DI at 40% under MTI is not beneficial for grain yield in relation to water consumption.

Biomass WUE showed significant variations across the three water regimes ($p < 0.05$). However, the type of irrigation did not significantly ($p > 0.05$) affect the biomass WUE in all the water regimes except at SDI at 40% ET_c where it was the highest. The DI significantly improved WUE by up to 45.8% and 21.2% for the 40% ET_c and 70% ET_c regimes, respectively. Therefore, in areas of water scarcity, cowpea can rather be grown for biomass than for grain yield. The cowpea variety used in this study (mixed brown) favours vegetative growth, thus gives more biomass than grain yield (Ilunga, 2014). The results of this study were consistent with those of Maleki et al. (2017), where the grain WUE was

highest at 80% of full irrigation compared to 60% and 40%. Similarly, Mousa and Qurashi (2017) reported increased WUE under water deficit imposed at various growth stages, except during a combination of the flowering and pod filling stages, where it decreased marginally. However, Ahmed and Suliman (2010) reported decreased WUE due to water deficit which was attributed to reduced photosynthetic activity.

The results reported in this study were consistent with those reported in other studies on MTI. For instance, Zhang et al. (2017) found significantly lower summer maize yields in MTI compared to SDI. In the same study, the yield of winter wheat was higher under SDI than MTI, but was not significantly different. Further, WUE was not significantly different between SDI and MTI in both maize and wheat. In another study, Zhang et al. (2016b), found that SDI marginally increased WUE of summer maize compared to MTI due to the former having a 2% higher average soil moisture content over the growing season. Therefore, the crop performance under MTI and SDI is not significantly different.

CONCLUSION

The results of this study showed that growth parameters of cowpea, i.e., LAI and phenology (time to 50% flowering) showed significant variations under MTI and SDI. For both irrigation system types, water deficit reduced the LAI and hastened the time to 50% flowering. The time to flowering was generally shorter under MTI than SDI. During the summer trial, grain yield was not affected significantly by irrigation type. However, there was significant variation in biomass between MTI and SDI with the former recording lower values than the latter. Biomass during the winter trial was not significantly different between MTI and SDI. Water deficit significantly reduced the yield and biomass of cowpea during the summer trial, especially at 40% ET_c. During the winter trial, water deficit had a significant effect only at 40% ET_c. Therefore, the hypothesis that the response of cowpea under both SDI and MTI is the same is rejected.

The grain WUE was improved by water deficit under SDI but only at 70% ET_c under MTI. Biomass WUE was significantly improved by increasing water deficit under both SDI and MTI, but this occurred to a greater extent under SDI. Therefore, the hypothesis that DI improves WUE of cowpea was accepted for biomass. The mixed brown variety of cowpea used in this study is highly vegetative and thus more suitable for biomass production rather than grain yield. Therefore, it is best suited as a leafy vegetable and fodder for human and animal consumption, respectively. This implies that DI could be a successful agricultural water management strategy in water-scarce regions.

Table 7. Water use efficiency for cowpea under MTI and SDI for summer experiment

Water regime	Amount of water applied (m ³ ·ha ⁻¹)	Water use efficiency (kg·m ⁻³)	
		Grain (SD)	Biomass (SD)
Moisture 100% ET _c	3 480	0.916 (0.182) ^{ab}	2.664 (0.358) ^a
SDI 100% ET _c	3 690	0.820 (0.188) ^{ab}	2.623 (0.298) ^a
Moisture 70% ET _c	2 520	0.954 (0.243) ^b	3.179 (0.416) ^b
SDI 70% ET _c	2 710	0.961 (0.259) ^b	3.170 (0.494) ^b
Moisture 40% ET _c	1 750	0.790 (0.369) ^a	3.519 (0.411) ^c
SDI 40% ET _c	1 620	0.860 (0.264) ^{ab}	3.825 (0.722) ^d
LSD ET _c)		0.093	0.169
LSD (Irrigation)		0.076	0.138
LSD (Irrigation x ET _c)		0.132	0.239
CV (%)		19.2	14.8

Mean values in same column followed by same superscript letter do not significantly differ at 5% significance level by LSD. Data in parentheses are the standard deviations.

ACKNOWLEDGEMENTS

The authors are grateful to the Association of African Universities (AAU), University of KwaZulu-Natal, Masinde Muliro University of Science and Technology and National Research Fund (NRF) of Kenya for funding this study.

REFERENCES

- ABAYOMI YA and ABIDOYE TO (2009) Evaluation of cowpea genotypes for soil moisture stress tolerance under screen house conditions. *Afr. J. Plant Sci.* **3** (10) 229–237.
- ABDOUL KARIM TD, SANOUSSI A, MAÂROUHI IM, FALALOU H and YACCOUBOU B (2018) Effect of water deficit at different stages of development on the yield components of cowpea (*Vigna unguiculata* L. Walp) genotype. *Afr. J. Biotechnol.* **17** (9) 279–287. <https://doi.org/10.5897/AJB2017.16347>
- AHMED FE and SULIMAN ASH (2010) Effect of water stress applied at different stages of growth on seed yield and water-use efficiency of cowpea. *Agric. Biol. J. N. Am.* **1** (4) 534–540.
- ALBAJI M, NASAB SB, BEHZAD M, NASERI A, SHAHNAZARI A, MESKARBASHEE M, JUDY F, JOVZI M and SHOKOOHFAR AR (2011) Water productivity and water use efficiency of sunflower under conventional and limited irrigation. *J. Food, Agric. Environ.* **9** (1) 202–209.
- ALLEN RG, PEREIRA LS, RAES D and SMITH M (1998) Crop evapotranspiration – Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. FAO, Rome, Italy.
- ANYIA AO and HERZOG H (2004) Water-use efficiency, leaf area and leaf gas exchange of cowpeas under mid-season drought. *Eur. J. Agron.* **20** (4) 327–339. [https://doi.org/10.1016/S1161-0301\(03\)00038-8](https://doi.org/10.1016/S1161-0301(03)00038-8)
- AYARS JE, PHENE CJ, HUTMACHER RB, DAVIS KR, SCHONEMAN RA, VAIL SS and MEAD RM (1999) Subsurface drip irrigation for row crops: a review of 15 years of research at the Water Management Research Laboratory. *Agric. Water Manage.* **42** 1–27. [https://doi.org/10.1016/S0378-3774\(99\)00025-6](https://doi.org/10.1016/S0378-3774(99)00025-6)
- BADIANE FA, DIOUF D, SANÉ D, DIOUF O, GOUDIABY V and DIALLO N (2004) Screening cowpea [*Vigna unguiculata* (L.) Walp.] varieties by inducing water deficit and RAPD analyses. *Afr. J. Biotechnol.* **3** (3) 174–178. <https://doi.org/10.5897/AJB2004.000-2030>
- BASARAN U, AYAN I, ACAR Z, MUT H and ASCI OO (2011) Seed yield and agronomic parameters of cowpea (*Vigna unguiculata* L.) genotypes grown in the Black Sea region of Turkey. *Afr. J. Biotechnol.* **10** (62) 13461–13464. <https://doi.org/10.5897/AJB11.2489>
- CISSE N (2001) Genotype x row spacing and environment interaction of cowpea in semi-arid zones. *Afr. Crop Sci. J.* **9** (2) 359–367. <https://doi.org/10.4314/acsj.v9i2.27606>
- DADSON RB, HASHEM FM, JAVAID I, JOSHI J, ALLEN AL and DEVINE TE (2005) Effect of water stress on the yield of cowpea (*Vigna unguiculata* L. Walp.) genotypes in the Delmarva Region of the United States. *J. Agron. Crop Sci.* **191** 210–217. <https://doi.org/10.1111/j.1439-037X.2005.00155.x>
- DAFF (Department of Agriculture, Forestry and Fisheries, South Africa) (2014) Production guidelines for cowpeas. Department of Agriculture, Forestry and Fisheries, Pretoria, South Africa.
- EHLERS JD and HALL AE (1997) Cowpea (*Vigna unguiculata* L. Walp.). *Field Crops Research* **53** 187–204. [https://doi.org/10.1016/S0378-4290\(97\)00031-2](https://doi.org/10.1016/S0378-4290(97)00031-2)
- FALOYE OT and ALATISE MO (2017) Response of soil moisture content, evapotranspiration, and yield of cowpea to varying water application. *Agric. Eng. Int. CIGR J.* **19** (4) 66–75.
- FAROOQ M, HUSSAIN M, WAHID A and SIDDIQUE KHM (2012) Drought stress in plants: an overview. In Aroca RS (ed) *Plant Responses to Drought Stress: From Morphological to Molecular Features*. Springer-Verlag, Berlin. 1–33. https://doi.org/10.1007/978-3-642-32653-0_1
- FATHI A and TARI DB (2016) Effect of drought stress and its mechanism in plants. *Int. J. Life Sci.* **10** 1–6. <https://doi.org/10.3126/ijls.v10i1.14509>
- FERERES E and SORIANO MA (2007) Deficit irrigation for reducing agricultural water use. *J. Exp. Bot.* **58** (2) 147–159. <https://doi.org/10.1093/jxb/erl165>
- FIGUEIREDO MVB, BEZERRA-NETO E and BURITY HA (2001) Water stress response on the enzymatic activity in cowpea nodules. *Braz. J. Microbiol.* **32** 195–200. <https://doi.org/10.1590/S1517-83822001000300006>
- HAMIDOU F, ZOMBRE G and BRACONNIER S (2007) Physiological and biochemical responses of cowpea genotypes to water stress under glasshouse and field Conditions. *J. Agron. Crop Sci.* **193** 229–237. <https://doi.org/10.1111/j.1439-037X.2007.00253.x>
- ILUNGA K (2014) Physiological responses of cowpea (*Vigna unguiculata*) to water stress under varying water regimes. Dissertation, University of KwaZulu-Natal.
- ISHIYAKU MF, SINGH BB and CRAUFURD PQ (2005) Inheritance of time to flowering in cowpea (*Vigna unguiculata* (L.) Walp.). *Euphytica* **142** (3) 291–300. <https://doi.org/10.1007/s10681-005-2435-0>
- KANDA EK, MABHAUDHI T and SENZANJE A (2018) Hydraulic and clogging characteristics of Moistube irrigation as influenced by water quality. *J. Water Suppl. Res. Technol.* **67** (5) 438–446. <https://doi.org/10.2166/aqua.2018.166>
- LYU W, NIU W, GU J, LI Y, ZOU X and ZHANG R (2016) Effects of moisture depth and density on tomato yield and quality in solar greenhouse. *Chin. J. Eco-Agric.* **24** (12) 1663–1673.
- MALEKI A, ESNAASHAR N and BIDABADI AA (2017) Impact of deficit irrigation on yield components, water use efficiency and yield response factor of cowpea in Khorramabad Iran. *J. Eng. Appl. Sci.* **12** (9) 2471–2479.
- MOUSA MAA and QURASHI ADA (2017) Growth and yield of cowpea (*Vigna unguiculata* L.) cultivars under water deficit at different growth stages. *Legume Res.* **40** (6) 1–8. <https://doi.org/10.18805/LR-384>
- NTOMBELA Z (2012) Growth and yield responses of cowpeas (*Vigna unguiculata* L.) to water stress and defoliation. Thesis, University of KwaZulu-Natal.
- PEKSEN E (2007) Yield performance of cowpea cultivars (*Vigna unguiculata* (L.) Walp) under rainfed and irrigated conditions. *Inter. J. Agric. Res.* **2** (4) 391–396. <https://doi.org/10.3923/ijar.2007.391.396>
- PRASAD PVV, STAGGENBORG SA and RISTIC Z (2008) Impacts of drought and/or heat stress on physiological, developmental, growth, and yield processes of crop plants. In: Ahuja LR, VR Reddy, SA Saseendran and Q Yus (eds) *Advances in Agricultural Systems Modeling, Response of Crops to Limited Water: Understanding and Modeling Water Stress Effects on Plant Growth Processes*. American Society of Agronomy, Crop Science Society of America and Soil Science Society of America, Wisconsin, USA. 301–355.
- REINDERS F, VAN DER STOEP I, LECLER N, GREAVES K, VAHRMEIJER J, BENADÉ N, DU PLESSIS F, VAN HEERDEN P, STEYN J, GROVÉ B, JUMMAN A and ASCOUGH G (2010) Standards and guidelines for improved efficiency of irrigation water use from dam wall release to root zone application. WRC Report No. K5/1482/4. Water Research Commission, Pretoria.
- SEBETHA ET, AYODELE VI, KUTU FR and MARIGA IK (2010) Yields and protein content of two cowpea varieties grown under different production practices in Limpopo province, South Africa. *Afr. J. Biotechnol.* **9** (5) 628–634. <https://doi.org/10.5897/AJB09.1132>
- SHIRINGANI RP and SHIMELIS HA (2011) Yield response and stability among cowpea genotypes at three planting dates and test environments. *Afr. J. Agric. Res.* **6** (14) 3259–3263.
- SINGH BB, AJEIGBE HA, TARAWALI SA, FERNANDEZ-RIVERA S and ABUBAKAR M (2003) Improving the production and utilization of cowpea as food and fodder. *Field Crops Res.* **84** 169–177. [https://doi.org/10.1016/S0378-4290\(03\)00148-5](https://doi.org/10.1016/S0378-4290(03)00148-5)
- SOUZA PJDOP, FARIAS VDS, LIMAMJA, RAMOSTF and SOUSA AML (2017) Cowpea leaf area, biomass production and productivity under different water regimes in Castanhal, Pará, Brazil. *Revista Caatinga* **30** (3) 748–759. <https://doi.org/10.1590/1983-21252017v30n323rc>
- SPRENT JI, ODEE DW and DAKORA FD (2009) African legumes: a vital but under-utilized resource. *J. Exp. Bot.* **61** (5) 1257–1265. <https://doi.org/10.1093/jxb/erp342>
- TIMKO MP and SINGH BB (2008) Cowpea, a multifunctional Legume. In: Moore PH and R Mings (eds) *Genomics of Tropical Crop Plants*. Springer, Berlin. 227–258. https://doi.org/10.1007/978-0-387-71219-2_10
- VAN RENSBURG WJ, VORSTER IHJ, VAN ZIJL JJB and VENTER SL (2007) Conservation of African leafy vegetables in South Africa. *Afr. J. Food Agric. Nutr. Dev.* **7** (3&4) 92–98.

- XUE W, NIU W, ZHANG Z and ZHANG K (2013) Effects of the tomato growth and water use efficiency in sunlight greenhouse by Moistube-irrigation. *Agric. Res. Arid Areas* **25** (6) 61–66.
- YANG W, TIAN L, DU T, DING R and YANG Q (2008) Research prospect of the water-saving irrigation by semi-permeable film. *J. Water Resour. Water Eng.* **19** (6) 60–63.
- YAO F, LIU H, LI Y, LIU F and YUAN N (2014) Research on ecophysiological parameters of navel orange under Moistube-irrigation. *J. Nanchang College Water Conserv. Hydroelec. Power* **33** (6) 11–14.
- YIN Y, SHEN L, GUO Y and ZHANG C (2017) The effect of different Moistube spacing of alternate micro-irrigation on water spinach growth in greenhouse. *Water Saving Irrig.* **42** (9) 1–4.
- ZHANG G, SHEN L and GUO Y (2016a) Experimental study on the growth status of cabbages in green house with Moistube irrigation technology. *Water Saving Irrig.* **41** (7) 6–8.
- ZHANG J, NIU W, ZHANG L and SHI L (2012) Experimental study on characters of wetted soil in moistube irrigation. *Sci. Soil Water Conserv.* **10** (6) 32–38.
- ZHANG M, NIU W, LU Z, LI Y, WANG J and QIU X (2017) Effect of Moistube-irrigation on crop yield and water use efficiency. *Chin. J. Eco-Agric.* **25** (11) 1671–1683.
- ZHANG M, NIU W, XU J and LI Y (2016b) Influences of micro-irrigation and subsoiling before planting on enzyme activity in soil rhizosphere and summer maize yield. *Chin. J. Appl. Ecol.* **27** (6) 1925–1934.
-